



**FIFRA 2(ee)  
Recommendation**

EPA Reg. No. 279-3588

**FOR DISTRIBUTION AND USE  
ONLY IN THE  
FOLLOWING STATES: AZ & CA**

THIS RECOMMENDATION, WHICH CONTAINS  
ADDITIONAL DIRECTIONS FOR USE, IS MADE AS  
PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS  
NOT BEEN SUBMITTED TO OR APPROVED BY THE  
US EPA.

This recommendation for Rhyme™ Fungicide is valid until December 31, 2027, or until withdrawn, canceled or suspended.

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.**

**THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**

**FOR THE SUPPRESSION OF *FUSARIUM SP.* VIA DRIP CHEMIGATION IN TOMATOES**

**Directions For Use**

Apply as directed through pressurized irrigation methods.

Crop	Disease	Rate	Application Directions	Retreatment interval (days)
Tomatoes (See label for listed crops)	<i>Fusarium oxysporum f. sp. lycopersici</i>  <i>Fusarium falciforme</i>	7 fl oz/A  (0.114 lbs ai/acre)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development at 2 weeks intervals. Best results have been obtained when Rhyme fungicide is applied preventatively, at 2 weeks after transplanting followed by subsequent applications at 2 weeks intervals.	14

**Restrictions**

- No single application may exceed 7 fl oz of product/A
- Do not apply more than 4 applications per year
- Do not apply more than 28 fl oz of product/A per year
- Do not apply more than 0.455 lb ai of flutriafol/A per year
- Restricted Entry Interval (REI) is 12 hours
- Pre harvest interval (PHI) – 0 days

R-4720 053122