

FIFRA 2(ee) Recommendation

EPA Reg. No. 279-3588

FOR DISTRIBUTION AND USE ONLY IN THE FOLLOWING STATES:

CA

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation for Rhyme ™ Fungicide is valid until January 15, 2028, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR MANAGEMENT OF GRAPE TRUNK DISEASE COMPLEX ON GRAPES AND TRUNK DISEASE COMPLEX ON TREE NUTS

DIRECTIONS FOR USE

For management of wood diseases on grapes and tree nuts, apply Rhyme fungicide as directed through pressurized irrigation methods such as macro-sprinkler, micro-sprinkler, or drip systems. For best results, use Rhyme fungicide as part of a long-term management program

Crop	Disease	Rate (fl oz/A)	Application Directions	RTI (Days)
Grapes	Vinewood diseases Botryosphaeria dothidea Eutypa lata Phomopsis canker Phaeoacremonium parasiticum Neofusicoccum parvum Neofusicoccum mediterraneum	5.0	New Vineyards (1-3 years old) Make initial application at the beginning of Spring Root Flush. Make second application post- harvest during Fall Root Flush. All applications must be made through a pressurized irrigation system. To achieve maximum performance, use a minimum of 10 fl oz/A for the year. Established Vineyards (>3 years old) Make initial application at the beginning of Spring Root Flush. Make second application 14 days later. Make third application at the start of Veraison. Make fourth application post-harvest during Fall Root Flush. All applications must be made through a pressurized irrigation system. To achieve maximum performance, use a minimum of 20 fl oz/A for the year	14
Tree Nuts (African Tree Nut, Almond, Brazil Nut, Burr Oak, Butternut, Cajou, Cashew, Castanha-Do-Maranhao, Coconut, Coquito Nut, Dika Nut, Guiana Chestnut, Hazelnut, Heartnut, Hickory Nut, Japanese Horse- Chestnut, Macadamia Nut, Monogongo Nut, Monkey-Pot, Pachira Nut, Pecan, Sapucaia Nut, and Walnut)	Trunk diseases Botryosphaeria dothidea Cytospora sorbicola Eutypa lata Ceratocystis destructans, Neofusicoccum parvum Calosphaeria pulchella, Phomopsis canker	7.0	Bearing Almond: Do not apply more than 2 applications per year. Make initial application of 7 fl oz/A at the beginning of Spring Root Flush. Make second application at Fall Root Flush. Apply through a pressurized irrigation system.	7
			Non- bearing almond and other tree nuts: Make initial application of 7 fl oz/A at the beginning of Spring Root Flush. Reapply at 30 day intervals as needed. Make final application at Fall Root Flush. Apply through a pressurized irrigation system. To achieve maximum performance, make 4 applications at 7 oz/A for the year	30

RESTRICTIONS:

Grapes

- No single application may exceed 5 fl oz of product/A.
- Do not apply more than 6 applications per year.
- Do not apply more than 30 fl oz of product/A per year.
- Do not apply more than 0.488 lb ai flutriafol/A per year.
- Restricted Entry Interval (REI) is 5 days for girdling or turning of grapes. For all other activities the REI is 12 hours.
- Preharvest Interval (PHI) Do not apply within 14 days of harvest

Tree Nuts

- No single application may exceed 7 fl oz of product/A.
- Do not apply more than 4 applications per year.
- Do not apply more than 28 fl oz of product/A per year.
- Do not apply more than 0.455 lb ai flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours
- Preharvest Interval (PHI) Do not apply within 14 days of harvest

R-4470 011823