



For use on almond; walnut; celery, Brassica leafy vegetables crop group 5; grapes (fresh, juice, table, wine, raisin); leafy vegetables, except Brassica, crop group 4; stone fruit; strawberry; tomatoes; vegetables, cucurbit, group 9; vegetables, fruiting, group 8-10; hops, peanut; pome fruit (apple, crabapple, loquat, Mayhaw, pear, oriental pear, quince); pecan and other tree nuts.

EPA Reg. No. 279-3588	EPA Est. 70815-GA-001
ACTIVE INGREDIENT:	By Wt.
*Flutriafol.....	22.7%
Other Ingredients.....	77.3%
Total:	100.0%

*Contains 2.08 pounds per gallon of the active ingredient flutriafol. Suspension concentrate.

**KEEP OUT OF REACH OF CHILDREN
WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**See additional precautionary statements and Directions for Use in booklet.
See First Aid statements on back panel.**

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.	
Note to Physician: No specific antidote. Treat symptomatically.	
For SPILLS Contact CHEMTREC 1-800-424-9300	

Sold By

 FMC Corporation
 2929 Walnut Street
 Philadelphia, PA 19104

Net Contents: 2.5 Gallons

SL-4457-1 083123 07-06-21

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Warning. Causes substantial but temporary eye injury. Harmful if swallowed. **DO NOT** get in eyes, or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators must wear: long-sleeved shirt and long pants, shoes plus socks, chemical-resistant gloves (Barrier Laminate, Butyl Rubber \geq 14 mils, Nitrile rubber \geq 14 mils, Neoprene rubber \geq 14 mils, Natural Rubber \geq 14 mils, Polyethylene, Polyvinyl chloride \geq 14 mils, or Viton® \geq 14 mils), and protective eyewear.

Mixers and Loaders must wear: long-sleeved shirt and long pants, shoes plus socks, and chemical resistant gloves (Barrier Laminate, Butyl Rubber \geq 14 mils, Nitrile rubber \geq 14 mils, Neoprene rubber \geq 14 mils, Natural Rubber \geq 14 mils, Polyethylene, Polyvinyl chloride \geq 14 mils, or Viton® \geq 14 mils) and protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This formulated product is toxic to fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Flutriafol has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in any manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

In California, this product may only be applied to fields that are not irrigated, or are irrigated using pressurized irrigation methods such as macro-sprinkler, micro-sprinkler, or drip systems.

Not for sale, distribution, or use in Nassau or Suffolk Counties of New York State.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Workers Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the Restricted Entry Interval (REI). The REI for each crop is located in the use directions for each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls, chemical resistant gloves, shoes plus socks, and protective eyewear.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store unused product in original container in a cool, dry, secure area.

Pesticide Disposal

Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable container equal to or less than 5 gallons

DO NOT reuse or refill this container.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure 2 more times. Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable container-greater than 5 gallons and less than 260 gallons

DO NOT reuse or refill this container.

Triple rinse as follows: Empty the remaining contents into application equipment or mix tank.

Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank and store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal.

Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PRODUCT DIRECTIONS

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH RESTRICTIONS, PRECAUTIONARY STATEMENTS, AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Not for use in greenhouses.

Product can be applied for all crops by foliar application or chemigation unless otherwise specified in this label.

Chemigation:

Application through Irrigation Systems (Chemigation):

Apply RHYME Fungicide through irrigation to crops at rates and timings specified in this label. Chemical tank and injector system must be thoroughly cleaned before and after use. Flush system with clean water.

Crop injury or lack of effectiveness can result from non-uniform distribution of RHYME Fungicide across the treated area. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Drip Irrigation: Use RHYME Fungicide for control of diseases at rates and timing as specified in this label. Ensure adequate soil moisture prior to utilizing RHYME Fungicide in a drip irrigation system. **DO NOT** apply RHYME Fungicide through drip irrigation that is deeper than 12" below the soil surface.

After application of RHYME Fungicide, continue to drip irrigate the area with water to flush the irrigation system. **DO NOT** allow RHYME Fungicide to remain in the irrigation system after the application is complete. The total volume of water, including the amount used for flushing the irrigation system, must be adequate to completely remove the product from the lines, but should be less than the amount that could oversaturate the beds (bed collapse can occur from over-saturation) in bedded production systems. Delay subsequent irrigation (water only) for at least 24 hours following drip application for best results.

RHYME fungicide Rate (fl. oz. /A)	Fluid ounces RHYME fungicide/1000 row feet based on planted row spacing												
	30" rows	34" rows	36" rows	38" rows	40" rows	44" rows	48" rows	60" rows	66" rows	72" rows	80" rows	84" rows	120" rows
5	0.29	0.33	0.34	0.36	0.38	0.42	0.46	0.57	0.63	0.69	0.77	0.80	1.15
7	0.40	0.46	0.48	0.51	0.54	0.59	0.64	0.80	0.88	0.96	1.07	1.12	1.61

Sprinkler Irrigation: Apply this product through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, micro and macro sprinkler or hand move irrigation systems. **DO NOT** apply this product through any other type of irrigation system except as specified on this label.

With Center Pivot irrigation equipment, use RHYME Fungicide only with equipment with drive systems that provide uniform water distribution. **DO NOT** use end guns for chemigation due to the non-uniform application pattern.

Add RHYME Fungicide to the pesticide supply tank containing sufficient water to maintain a continuous flow by the injection equipment. Maintain agitation during the entire application period.

For continuous-move irrigation systems, apply the labeled rate for that crop in ½ acre-inch or less per acre. For stationary or non-continuous moving systems, inject RHYME Fungicide spray mixture during the last 20-30 minutes of the set.

DO NOT apply when winds are greater than 10-15 mph to avoid drift or wind skips. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. Thorough coverage of foliage is required for good control.

Operating Requirements for Application through Irrigation Systems:

1. **DO NOT** use chemigation when conditions are favorable for drift to non-target areas.
2. To prevent water-source contamination from backflow, a functional check valve, vacuum relief valve, and low-pressure drain should be located on the irrigation pipeline.
3. To prevent backflow back toward the injection pump, the pesticide injection pipeline must be equipped with a functional, automatic, quick-closing check valve.
4. To prevent fluid from being withdrawn from the supply tank when the irrigation system is shut down, the pesticide injection pipeline should also be equipped with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock.
5. The system must also contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops and a functional pressure switch to stop the pump motor when water pressure decreases to a point at which pesticide distribution is adversely affected.
6. A metering pump, constructed of materials compatible with pesticides and capable of being fitted with a system interlock, such as a positive displacement injection pump (e.g., a diaphragm pump), must be included in the system.
7. A knowledgeable person responsible for the chemigation system should shut the system down and turn the irrigation water off, ensuring enough time for the pesticide to be flushed through all lines and nozzles.
8. No irrigation system, including those in greenhouses, used to distribute pesticides can be connected to a public water source unless safety measures and devices prescribed in the pesticide label for such connection are in place.

Specific Instructions for Public Water Systems:

1. Public water system means a system that provides piped water for human consumption if the system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. All measures and devices listed in the above section, "Operating Requirements for Application through Irrigation Systems", must be operational for connection to a public water system.
3. Additionally, chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There must be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

Rainfastness: RHYME Fungicide is rainfast 2 hours after application. Disease control will be reduced if rainfall occurs within 2 hours of application.

Crop Rotation: Fields treated with RHYME Fungicide may be planted to the following crops and plant back intervals.

Crop	Plant back interval
Alfalfa, barley, field corn; field corn grown for seed; popcorn; sweet corn; cotton; cucurbits; fruiting vegetables; grapes; hops; Brassica leafy vegetables crop group 5; leafy vegetables, except Brassica, crop group 4; peanut; pome fruit; grain sorghum; soybean; stone fruit; strawberry; sugar beet; tomato; tree nuts; triticale; or wheat	Immediately
Rice	30 days
All other crops	Prohibited

RATE EQUIVALENCY TABLE

Product Rate (fl. oz./A)	Equivalent Active Ingredient Rate (lb a.i./A)
3.5	0.057
4	0.065
5	0.081
6	0.098
6.5	0.106
7	0.114
8	0.130
12	0.20
15	0.24
16	0.26
24	0.39

Use of Adjuvants: An adjuvant may be used with RHYME Fungicide unless otherwise specified in the crop use directions. RHYME Fungicide is compatible with most products, however not all have been tested. Use the compatibility jar test to ensure physical compatibility.

Tank Mixing: RHYME Fungicide may be tank mixed with other fungicides, herbicides, insecticides, and/or other additives unless prohibited on the label of the tank mix partner. Follow more restrictive labeling of any tank mix partner. Although RHYME Fungicide is compatible with most products, not all combinations have been tested. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Use the compatibility jar test to ensure physical compatibility. Before applying any tank mixture not specifically recommended on this label, the crop safety of the target crop should be confirmed by applying the mixture to a small area of the target crop in accordance to the label instructions.

Compatibility Jar Test: RHYME Fungicide is compatible with most products, however not all have been tested. Use the following compatibility test to ensure physical compatibility.

Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add wettable powders and water dispersible granular products first, next liquid flowables, then emulsifiable concentrates, and last liquid soluble products. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Once compatibility has been proven, use the same procedure for adding products to the spray tank. When tank mixing, follow more restrictive labeling of any tank mix partner. **DO NOT** tank mix with any product that contains a prohibition on tank mixing.

Spray Equipment/Volume: Apply RHYME Fungicide in sufficient water volume to ensure thorough coverage as a broadcast foliar application for good disease control.

For ground application, use a minimum of 30 gallons of spray solution per acre by ground sprayer for hops, and 10 gallons of spray solution per acre by ground sprayer for all other crops. For aerial application, use aircraft spray equipment in a minimum of 10 gallons of spray solution per acre for tree nut crops and hops, 2 gallons of spray solution per acre for corn, cotton, grain sorghum, soybean, wheat and triticale, and 5 gallons of spray solution per acre for all other crops.

Resistance Management: For resistance management, RHYME Fungicide contains a Group 3 fungicide (Flutriafol.)

Any fungal population may contain individuals naturally resistant to RHYME Fungicide and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of RHYME Fungicide or other Group 3 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact your pesticide distributor or university extension specialist to report resistance.

High Disease Pressure: Where a range is given, use the higher specified rate of RHYME Fungicide and/or reduced spray interval when disease pressure is high and/or conditions are favorable for disease development.

CROP USE DIRECTIONS

Use Directions for Almond

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Brown Rot Blossom Blight (<i>Monilinia</i> spp.) Alternaria Leaf and Fruit Spot (<i>Alternaria alternata</i>) Anthracnose (<i>Colletotrichum acutatum</i>) Scab (<i>Cladosporium carpophilum</i>) Shot Hole (<i>Wilsonomyces carpophilus</i>) Leaf Rust (<i>Tranzschelia discolor</i>)	7 (0.114 lb ai/A)	For Brown Rot Blossom Blight start applications at 5-10% bloom followed by an application at 50-100% bloom. For all other diseases apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	7
Restrictions: <ul style="list-style-type: none"> No single application may exceed 7 fl oz of product (0.114 lb ai)/A. DO NOT apply more than 4 applications per year. DO NOT apply more than 28 fl oz of product (0.455 lb ai)/A per year. DO NOT apply more than 0.455 lb ai of flutriafol/A per year. Restricted Entry Interval (REI) is 12 hours. Preharvest Interval (PHI): DO NOT apply within 14 days of harvest. 			

Use Directions for Walnut including black and English (Persian) walnuts

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Not for Use in CA Botryosphaeria panicle and shoot blight Not for Use in CA Phomopsis canker	7 (0.114 lb ai/A)	Apply preventatively and repeat as necessary on a 7-day interval when conditions are favorable for disease development.	7
Restrictions: <ul style="list-style-type: none"> No single application may exceed 7 fl oz of product (0.114 lb ai)/A. The minimum retreatment interval is 7 days. DO NOT apply more than 4 applications per year. DO NOT apply more than 28 fl oz of product (0.455 lb ai)/A per year. DO NOT apply more than 0.455 lb ai of flutriafol/A per year. Restricted Entry Interval (REI) is 12 hours. Preharvest Interval (PHI): DO NOT apply within 14 days of harvest. 			

Use Directions for Brassica Leafy Vegetables Crop Group 5.

See below for a list of crops.

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Black Spot (<i>Alternaria</i> spp.) Cercospora Leaf Spot (<i>Cercospora</i> spp.) Powdery Mildew (<i>Erysiphe polygoni</i>)	5 – 7 (0.081 – 0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	7
List of Brassica Leafy Vegetables Crop Group 5: broccoli; broccoli Chinese (gai lon); broccoli raab (rapini); Brussels sprouts; cabbage; cabbage, Chinese (bok choy); cabbage Chinese, (napa); cabbage, Chinese mustard (gai choy); cauliflower; cavalo broccolo; collards; kale; kohlrabi; mizuna; mustard greens; mustard spinach; rape greens and cultivars and/or hybrids of these crops.			
Restrictions: <ul style="list-style-type: none"> No single application may exceed 7 fl oz of product (0.114 lb ai)/A. DO NOT apply more than 4 applications per year. DO NOT apply more than 28 fl oz of product (0.455 lb ai)/A per year. DO NOT apply more than 0.455 lb ai of flutriafol/A per year. Restricted Entry Interval (REI) is 12 hours. Preharvest Interval (PHI): DO NOT apply within 7 days of harvest. 			

Use Directions for Celery and Chinese Celery

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Alternaria Leaf Spot (<i>Alternaria sonchi</i> , <i>A. spp.</i>) Anthracnose (<i>Colletotrichum dematium</i>) Early Blight (<i>Cercospora apii</i>) Late Blight (<i>Septoria apiicola</i>)	5 – 7 (0.081 – 0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	7
Fusarium Yellows/ Fusarium Wilt (Suppression only) (<i>Fusarium oxysporum f.sp apii</i>)	5 – 7 (0.081-0.114 lb ai/A)	Product is to be applied through drip irrigation. Recommend initial application at transplanting. For spring and fall planting, a second application is required, approximately 30 days later. For fall planting, additional applications may be required at 14 day intervals, for a maximum of 4 applications per season.	14
Restrictions: <ul style="list-style-type: none"> No single application may exceed 7 fl oz of product (0.114 lb ai)/A. DO NOT apply more than 4 applications per year. DO NOT apply more than 28 fl oz of product (0.455 lb ai)/A per year. DO NOT apply more than 0.455 lb ai of flutriafol/A per year. Restricted Entry Interval (REI): The REI is 12 hours. Preharvest Interval (PHI): DO NOT apply within 7 days of harvest. 			

Use Directions for Grapes (fresh, juice, table, wine, raisin)

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Powdery Mildew (<i>Uncinula necator</i>)	4 – 5 (0.065-0.081 lb ai/A)	Foliar Application: Begin application at 6 to 10 inch shoot growth.	14 – 21
	5 (0.081 lb ai/A)	Drip Irrigation Application: Apply preventively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development and to provide extended foliar protection. Best results have been obtained when RHYME fungicide is applied preventively, as early as bud break, to ensure time for movement of flutriafol into the foliage. Once fruit are present, foliar applications are needed for fruit protection.	14
Black Rot (<i>Guignardia bidwellii</i>)	4 – 5 (0.065-0.081 lb ai/A)	Foliar Application: Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	14 – 21
Restrictions: <ul style="list-style-type: none"> No single application may exceed 5 fl oz of product (0.081 lb ai)/A. The minimum retreatment interval is 14 days. DO NOT apply more than 6 applications per year. DO NOT apply more than 30 fl oz of product (0.488 lb ai)/A per year. DO NOT apply more than 0.488 lb ai of flutriafol/A per year. Restricted Entry Interval (REI): For foliar applications: The REI is 5 days for girdling or turning of grapes. For soil-applied applications: The REI is 12 hours for girdling or turning of grapes. The REI for all other activities is 12 hours. Preharvest Interval (PHI): DO NOT apply within 14 days of harvest. 			

Use Directions for Leafy Vegetables, except Brassica, Crop Group 4.
See below for a list of crops.

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Alternaria Leaf Spot <i>(Alternaria sonchi,</i> <i>A. spp.)</i> Anthracnose <i>(Microdochium</i> <i>panattonianum,</i> <i>Colletotrichum</i> <i>dematium)</i> Cercospora Leaf Spot <i>(Cercospora spp.)</i> Powdery Mildew <i>(Erysiphe</i> <i>cichoracearum,</i> <i>Leveillula taurica)</i> Septoria Leaf Spot <i>(Septoria spp)</i>	5 – 7 (0.081 – 0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development.	7
List of Leafy Vegetables, except Brassica, Crop Group 4: amaranth (Chinese spinach); amaranth (leafy); amaranth (tampala); arugula (roquette); cardoon; celtuce; chervil; chrysanthemum (edible-leaved); chrysanthemum (garland); corn salad; cress (garden); cress (upland); dandelion (leaves); dock (sorrel); endive (escarole); fennel, Florence; lettuce (head and leaf); orach; parsley; purslane (garden); purslane (winter); radicchio (red chicory); rhubarb; spinach; spinach (New Zealand); spinach (vine); Swiss chard and cultivars and/or hybrids of these crops.			
Restrictions: <ul style="list-style-type: none"> No single application may exceed 7 fl oz of product (0.114 lb ai)/A. DO NOT apply more than 4 applications per year. DO NOT apply more than 28 fl oz of product (0.455 lb ai)/A per year. DO NOT apply more than 0.455 lb ai of flutriafol/A per year. Restricted Entry Interval (REI): The REI is 12 hours. Preharvest Interval (PHI): DO NOT apply within 7 days of harvest. 			

Use Directions for Stone Fruit.
See below for a list of crops.

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Powdery Mildew <i>(Sphaerotheca</i> <i>pannosa)</i>	7 (0.114 lb ai/A)	Start applications at petal fall. Repeat as necessary if disease conditions are favorable.	7
Not for Use in CA Rusty Spot <i>(Podosphaera</i> <i>leucotricha)</i>	7 (0.114 lb ai/A)	Start applications at petal fall and continue through 3rd Cover (3C) or until terminal shoot growth ceases and shoots harden off or conditions are no longer favorable for disease development.	7
Brown Rot Blossom Blight <i>(Monilinia spp.)</i>	7 (0.114 lb ai/A)	Start applications at 1-5% bloom followed by an application at 50-100% bloom.	7
Fruit Brown Rot <i>(Monilinia spp.)</i>	7 (0.114 lb ai/A)	Apply one to two weeks prior to harvest.	7
Shot Hole <i>(Wilsonomyces</i> <i>carophilus)</i>	7 (0.114 lb ai/A)	Apply at 50-100% bloom. Repeat as necessary if disease conditions are favorable for disease development.	7
Cherry Leaf Spot <i>(Blumeriella jaapii)</i>	7 (0.114 lb ai/A)	Start applications at petal fall. Repeat as necessary if disease conditions are favorable for disease development.	7
List of Stone Fruit: Apricot, Sweet Cherry, Tart Cherry, Nectarine, Peach, Plum, Chickshaw Plum, Damson Plum, Japanese Plum, Plumcot, Prune.			
Restrictions: <ul style="list-style-type: none"> No single application may exceed 7 fl oz of product (0.114 lb ai)/A. DO NOT apply more than 4 applications per year. DO NOT apply more than 28 fl oz of product (0.455 lb ai)/A per year. DO NOT apply more than 0.455 lb ai of flutriafol/A per year. Restricted Entry Interval (REI) is 12 hours. Preharvest Interval (PHI): DO NOT apply within 7 days of harvest. 			

Use Directions for Strawberry

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Powdery Mildew (<i>Podosphaera aphanis</i>)	5 – 7 (0.081-0.114 lb ai/A)	Foliar Application: Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	7
Not for Use in CA Leaf Rust (<i>Phragmidium potentillae</i>) Leaf Spot (<i>Mycosphaerella fragariae</i>)	7 (0.114 lb ai/A)	Drip Irrigation: Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development. Best results for both foliar and soilborne pathogens have been obtained when RHYME fungicide is applied preventively, shortly after transplanting, to provide protection of the roots from soilborne pathogens and to ensure time for movement of flutriafol into the foliage for protection against foliar pathogens. Once fruit are present, foliar applications are needed for fruit protection.	14
Not for Use in CA Anthracnose (<i>Colletotrichum acutatum</i>)	7	Drip Irrigation: Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development. Best results for both foliar and soilborne pathogens have been obtained when RHYME fungicide is applied preventively, shortly after transplanting, to provide protection of the roots from soilborne pathogens and to ensure time for movement of flutriafol into the foliage for protection against foliar pathogens. Once fruit are present, foliar applications are needed for fruit protection.	14
Not for use in CA Charcoal Rot (<i>Macrophomina phaseolina</i>)	7 (0.114 lb ai/A)	Product is to be applied through drip irrigation (see Chemigation for directions). Recommend initial application at transplanting. For spring and fall planting sequential applications are recommended on a 14 day interval.	14
Restrictions: <ul style="list-style-type: none"> No single application may exceed 7 fl oz of product (0.114 lb ai)/A. The minimum retreatment interval is 7 days. DO NOT apply more than 4 applications per year. DO NOT apply more than 28 fl oz of product (0.228 lb ai)/A per year. DO NOT apply more than 0.455 lb ai of flutriafol/A per year. Restricted Entry Interval (REI) is 12 hours. Preharvest Interval (PHI): May be applied the day of harvest (0 day PHI). 			

Use Directions for Tomatoes.
See below for a list of crops.

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Powdery Mildew (<i>Leveillula Taurica</i> , <i>Oidium neolycopersici</i>) Anthracnose (<i>Collectotrichum coccodes</i>) Target Spot (<i>Corynespora cassiicola</i>) Not for Use in CA Septoria Leaf Spot (<i>Septoria lycopersici</i>) Not for Use in CA Cercospora Leaf Spot (<i>Cercospora capsici</i>)	Foliar: 5 – 7 (0.081-0.114 lb ai/A) Drip: 7 (0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	Foliar – 7 Drip - 14
Black Mold (<i>Alternaria alternata</i>) Early Blight (<i>Alternaria solani</i>)	3.5 – 7 (0.057-0.114 lb ai/A)	Tank mix with mancozeb for improved disease control. Lower rate can be used when tank mixed with mancozeb or other effective fungicides.	7
Not for use in CA Charcoal Rot (<i>Macrophomina phaseolina</i>)	7 (0.114 lb ai/A)	Apply through drip irrigation (see Chemigation for directions). Recommend initial application at transplanting. A second application is required approximately 14 days later. Two additional applications may be applied if necessary.	14
Not for use in CA Fusarium Wilt (Suppression) (<i>Fusarium oxysporum f. sp lycopersici</i>)	7 (0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development. Best results for soilborne pathogens has been obtained when RHYME fungicide is applied preventively, shortly after transplanting, followed by subsequent applications at 14 day intervals to provide protection of the roots from soilborne pathogens.	14
List of Tomatoes: Bush Tomato, Cocona, Currant Tomato, Garden Huckleberry, Goji Berry, Groundcherry, Narnajilla, Sunberry, Tomatillo, and Tree Tomato and cultivars, varieties and/or hybrids of these crops.			
Restrictions:			
<ul style="list-style-type: none"> • No single application may exceed 7 fl oz of product (0.114 lb ai)/A. • The minimum retreatment interval is 7 days. • DO NOT apply more than 4 applications per year. • DO NOT apply more than 28 fl oz of product/A per year. • DO NOT apply more than 0.455 lb ai of flutriafol/A per year. • Restricted Entry Interval (REI) is 12 hours. • Preharvest Interval (PHI): May be applied the day of harvest (0 day PHI). When tank mixed with mancozeb fungicide, the PHI is 5 days. 			

Use Directions for Vegetables, Cucurbit, Group 9.
See below for a list of crops.

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Not for Use in CA Anthraxnose (<i>Colletotrichum</i> spp.) Powdery Mildew (<i>Sphaerotheca fuliginea</i> , <i>Erysiphe cichoracearum</i>) Not for Use in CA Cercospora Leaf Spot (<i>Cercospora citrulline</i>) Gummy Stem Blight (<i>Didymella bryoniae</i>) Target Spot (<i>Corynespora cassiicola</i>)	Foliar: 5 - 7 (0.081- 0.114 lb ai/A) Drip: 7 (0.114 lb ai/A)	Apply preventively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development and to provide extended foliar protection. Best results for both foliar and soilborne pathogens have been obtained when RHYME fungicide is applied preventively, as early as 14 days after transplanting, to provide protection of the roots from soilborne pathogens and to ensure time for movement of flutriafol into the foliage for protection against foliar pathogens. Once fruit are present, foliar applications are needed for fruit protection. For gummy stem blight and target spot, tank mix with mancozeb for resistance management and in fields with known triazole resistance, apply RHYME Fungicide with mancozeb or alternative effective fungicide mode of action for disease control.	Foliar – 7 Drip - 14
Charcoal Rot (<i>Macrophoma phaseolina</i>)	7 (0.114 lb ai/A)	Apply through drip irrigation (see Chemigation section). Make initial application at transplanting. Make a second application 14 days later. Additional applications may be made on a 14 day interval, as needed.	14
Suppression of: Fusarium Wilt (<i>Fusarium oxysporum</i>)	7 (0.114 lb ai/A)	Apply preventively or when conditions are favorable for disease development and repeat application every 14 days to provide extended in-season suppression. Best results for soilborne disease suppression have been obtained when RHYME fungicide is applied preventively, as early as 14 days after transplanting, to provide protection of the roots from soilborne pathogens. RHYME fungicide may be alternated with other non-Group 3 mode of action fungicides to improve Fusarium management.	14

List of Cucurbits:

Chayote (fruit); Chinese waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); *Momordica* spp (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); muskmelon (includes true cantelope, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon); pumpkin; squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon* and hybrids and/or varieties of (*Citrullus* spp., *Cucumis* melo, *Cucurbita pepo*, *Lagenaria* spp., *Luffa* spp. *Momordica* spp. *Sechium edule*).

Restrictions for all cucurbit group 9 uses:

- No single application may exceed 7 fl oz of product (0.114 lb ai)/A.
- **DO NOT** apply more than 4 applications per year.
- **DO NOT** apply more than 28 fl oz of product (0.455 lb ai)/A per year.
- **DO NOT** apply more than 0.455 lb ai of flutriafol/A per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Interval (PHI): May be applied the day of harvest (0 day PHI).

Additional Watermelon only Restrictions:

- **DO NOT** apply more than 2 sequential applications, including soil and foliar applications of any Group 3 fungicide, before rotating with a fungicide from a different FRAC group.
- Use nonionic surfactants (NIS) only.
- **DO NOT** use organo silicone adjuvants, or crop oil concentrate (COC), or methylated seed oil (MSO) surfactants.

*Precaution for watermelon use: Under certain environmental conditions, applications of RHYME Fungicide may result in phytotoxicity symptoms that may reduce watermelon yields. FMC recommends the grower and/or user test this product to determine its suitability for use in watermelon. FMC provides this product to the grower and/or user solely to the extent that the benefit and utility, in the sole opinion of the grower and/or user, outweighs the potential of crop phytotoxicity following applications of RHYME Fungicide to watermelon. The decision to use or not use RHYME Fungicide in watermelon must be made by each individual grower and/or user on the basis of possible crop phytotoxicity, the severity of disease incidence, the cost of alternative disease control options, and other factors.

Use Directions for Hops

Not for use in CA

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Powdery Mildew (<i>Podosphaera macularis</i> , <i>Sphaerotheca macularis</i>)	5-7 (0.081 – 0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development. Apply a minimum of 30 gallons per acre spray volume for ground applications and 10 gallons per acre spray volume for aerial applications.	14
Restrictions: <ul style="list-style-type: none"> No single application may exceed 7 fl oz product (0.114 lb ai)/A DO NOT apply more than 4 applications per year DO NOT apply more than 28 fl oz (0.455 lb ai)/A per year DO NOT apply more than 0.455 lb ai of flutriafol/A per year. An adjuvant may be used with RHYME Fungicide Restricted Entry Interval (REI) is 12 hours Preharvest Interval (PHI): DO NOT apply within 7 days of harvest 			

Use Directions for Peanut

Not for use in CA

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Early Leaf Spot (<i>Cercospora arachidicola</i>) Late Leaf Spot (<i>Cercosporidium personatum</i>) Peanut Rust (<i>Puccinia arachidis</i>) Leaf Scorch/ Pepper Spot (<i>Leptosphaerulina crassiasca</i>) Web Blotch (<i>Phoma arachidicola</i>)	5 – 7 (0.081 – 0.114 lb ai/A)	Apply on a preventive spray schedule for effective disease control in peanuts. RHYME Fungicide can be tank mixed with other fungicides for improved disease control, broader spectrum and resistance management.	14
Suppression of Soil-Borne Diseases: White Mold/Southern Stem Rot (<i>Sclerotium rolfsii</i>)	5 – 7 (0.081 – 0.114 lb ai/A)	For soil-borne diseases, rainfall or irrigation will optimize activity of RHYME Fungicide. Tank-mix with other labeled fungicides for improved activity on soil-borne diseases.	14
Restrictions: <ul style="list-style-type: none"> No single application may exceed 7 fl oz of product (0.114 lb ai)/A. DO NOT apply more than 4 applications per year. DO NOT apply more than 28 fl oz of product (0.455 lb ai)/A per year. DO NOT apply more than 0.455 lb ai of flutriafol/A per year. Restricted Entry Interval (REI) is 12 hours. Preharvest Interval (PHI): DO NOT apply within 7 days of harvest. 			

**Use Directions for Pecan and Other Tree Nuts.
See below for list of crops.**

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Pecan Scab (<i>Fusicladium effusum</i>) Powdery Mildew (<i>Microsphaera penicillata</i>) Alternaria Leaf and Fruit Spot (<i>Alternaria alternata</i>) Anthracnose (<i>Colletotrichum acutatum</i>) Downy Spot (<i>Mycosphaerella caryigena</i>) Zonate Leaf Spot (<i>Cristulariella moricola</i>)	5 – 7 (0.081 – 0.114 lb ai/A)	For Pecan Scab control, start application at bud break. For all other disease apply preventatively or when conditions remain favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	7
Eastern Filbert Blight (<i>Anisogramma anomalea</i>)	7	For Hazelnut only apply preventatively or when conditions remain favorable for disease development. Repeat as necessary if conditions are favorable for disease development.	7
List of Tree Nuts: African Tree Nut, Brazil Nut, Burr Oak, Butternut, Cajou, Cashew, Castanha-Do-Maranhao, Coconut, Coquito Nut, Dika nut, Guiana Chestnut, Hazelnut, Heartnut, Hickory Nut, Japanese Horse- Chestnut, Macadamia Nut, Monogongo Nut, Monkey-Pot, Pachira Nut, Pecan, Sapucaia Nut.			
Restrictions: <ul style="list-style-type: none"> • No single application may exceed 7 fl oz of product (0.114 lb ai)/A. • DO NOT apply more than 4 applications per year. • DO NOT apply more than 28 fl oz of product (0.455 lb ai)/A per year. • DO NOT apply more than 0.455 lb ai of flutriafol/A per year. • Restricted Entry Interval (REI) is 12 hours. • Preharvest Interval (PHI): DO NOT apply within 14 days of harvest. 			

Use Directions for Vegetables, Fruiting, Group 8-10.
See below for list of crops.

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Powdery Mildew <i>(Leveillula taurica)</i> Anthracnose <i>(Colletotrichum coccodes, C. acutatum)</i> Cercospora Leaf Spot <i>(Cercospora capsici)</i> Not for Use in CA Black Mold <i>(Alternaria alternata)</i> Not for Use in CA Early Blight <i>(Alternaria solani)</i>	7 (0.114 lb ai/A)	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development. Best results for both foliar and soilborne pathogens have been obtained when RHYME fungicide is applied preventively, shortly after transplanting, to provide protection of the roots from soilborne pathogens and to ensure time for movement of flutriafol into the foliage for protection against foliar pathogens. Once fruit are present, foliar applications are needed for fruit protection.	Foliar – 7 Drip Irrigation 14
Not for Use in CA Target Spot <i>(Corynespora cassiicola)</i>	7 (0.114 lb ai/A)	Only apply through drip irrigation. Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions remain favorable for disease development. Best results for both foliar and soilborne pathogens have been obtained when RHYME fungicide is applied preventively, shortly after transplanting, to provide protection of the roots from soilborne pathogens and to ensure time for movement of flutriafol into the foliage for protection against foliar pathogens. Once fruit are present, foliar applications are needed for fruit protection.	14
Charcoal Rot <i>(Macrophomina phaseolina)</i> Not for use in CA Fusarium Wilt. <i>(Fusarium oxysporum)</i>	7 (0.114 lb ai/A)	Product is to be applied through drip irrigation (see Chemigation section). Make initial application at transplanting. Make a second application 14 days later. Additional applications may be made on a 14 day interval, as needed.	14
List of Fruiting Vegetables: African Eggplant, Bell Pepper, Eggplant, Martynia, Non-Bell Pepper, Okra, Pea Eggplant, Pepino, Roselle, Scarlet Eggplant, and cultivars, varieties and/or hybrids of these crops.			
Restrictions: <ul style="list-style-type: none"> • No single application may exceed 7 fl oz of product (0.114 lb ai)/A. • DO NOT apply more than 4 applications per year. • DO NOT apply more than 28 fl oz of product (0.455 lb ai)/A per year. • DO NOT apply more than 0.455 lb ai of flutriafol/A per year. • Restricted Entry Interval (REI) is 12 hours. • Preharvest Interval (PHI): May be applied the day of harvest (0 day PHI). 			

Use Directions for Pome Fruit (Apple, Crabapple, Loquat, Mayhaw, Pear, Oriental Pear, and Quince)

Diseases	Product Rate (fl oz/A)	Application Directions	Application Interval (Days)
Powdery Mildew (<i>Podosphaera leucotricha</i>) Quince Rust (<i>Gymnosporangium clavipes</i>) Cedar Apple Rust (<i>Gymnosporangium juniperi-virginianae</i>) Not for Use in CA Brooks Fruit Spot (<i>Mycosphaerella pomii</i>) Not for Use in CA Flyspeck (<i>Zygophiala jamacaicensis</i>) (formerly <i>Schizothyrium pomii</i>)	4 – 6 (0.065 – 0.098 lb ai/A)	Initiate applications at green tip and continue through cover sprays.	10
Apple Scab (<i>Venturia inaequalis</i>) Not for Use in CA Sooty Blotch (<i>Gleodes pomigena</i>) Not for Use in CA Bot Rot or White Rot (<i>Botryosphaeria dothidea</i>) Not for Use in CA Black Rot (<i>Botryosphaeria obtusa</i>) Not for Use in CA Bitter Rot (<i>Colletotrichum gloeosporoides</i> , <i>C.acutatum</i>) Not for Use in CA Alternaria Blotch (<i>Alternaria mali</i>)	6.5 (0.106 lb ai/A)	Initiate applications at green tip or when environmental conditions are favorable for primary scab development. Continue applications through the duration of primary scab. Mix RHYME Fungicide with a protectant such as mancozeb or captan fungicide for scab control. Tank mix applications containing RHYME Fungicide with a protectant such as mancozeb or captan fungicide will also effectively control listed summer rots when applied at cover spray timings.	7 - 10
Pear Scab (<i>Venturia pirina</i>)	6.5 (0.106 lb ai/A)	Start application at Pink Bud and continue through the duration of the Primary Scab period. Mix RHYME Fungicide with a protectant such as mancozeb or captan fungicide for improved scab control.	7 - 10
<p>Restrictions:</p> <ul style="list-style-type: none"> • No single application may exceed 6.5 fl oz of product (0.106 lb ai)/A. • DO NOT apply more than 4 applications per year. • DO NOT apply more than 26 fl oz of product (0.423 lb ai)/A per year. • DO NOT apply more than 0.423 lb ai of flutriafol/A per year. • Restricted Entry Interval (REI) is 12 hours. • Preharvest Interval (PHI): DO NOT apply within 14 days of harvest. 			

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or FMC, and Buyer assumes the risk of any such use.

To the extent consistent with applicable law, FMC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Condition of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

RHYME is a trademark of The FMC Corporation or an affiliate.