



FIFRA 2(ee) Recommendation
EPA Reg. No. 279-3588
FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA
THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation for Rhyme™ fungicide is valid until December 31, 2027, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.
THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR SUPPRESSION OF FUSARIUM SP. VIA CHEMIGATION IN CELERY

Directions for Use
Rhyme fungicide will provide suppression of listed diseases.
Product is to be applied through drip irrigation. Recommend initial application at transplanting. For spring and fall planting, a second application is required, approximately 30 days later. For fall planting, additional applications may be required at 14 day intervals, for a maximum of 4 applications per season.

Crop	Pest	Rate (fl oz/A)	PHI	Retreatment Interval (Days)	Restrictions
Celery	Fusarium Yellows /Fusarium Wilt (Suppression only) (<i>Fusarium oxysporum f.sp apii</i>)	5-7	Do not apply within 7 days of harvest	14	<ul style="list-style-type: none"> Do not apply more than 7.0 fl. oz. (0.114 lb ai) product/A/ application. Do not apply more than 28 fl. oz. of product/A/year. Do not make more than 4 applications/ year. Do not apply more than 0.455 lb. ai of flutriafol or flutriafol containing products/A/year.

R-4875 010424 (Supercedes R-4442 040920)



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104