



FIFRA 2(EE) Recommendation
EPA Reg. No. 279-3588
FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA
THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(EE) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This 2EE Recommendation for Rhyme™ Fungicide is valid until December 31, 2029, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.
THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

For Management of Walnut Mold in Walnuts (including black and English (Persian) walnuts)

Directions for Use				
Apply RHYME Fungicide in sufficient water volume to ensure thorough coverage as a foliar application. Refer to University and/or extension guidelines for best application timing.				
Crop	Disease	Rate (Fl. Oz./A)	Application Directions	Retreatment Interval (Days)
Walnut	Walnut Mold caused by: <i>Botryosphaeria</i> <i>Phomopsis</i> <i>Fusarium</i> <i>Aspergillus niger</i>	7	Begin fungicide applications preventatively. Continue as needed on a 7- day interval or when conditions are favorable for disease development.	7
To maintain performance of RHYME Fungicide and other fungicides in the same FRAC 3 group, tank mix or rotate with a different fungicide group for good disease resistance management.				
ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED RHYME FUNGICIDE LABEL (EPA REG. NO. 279-3588) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.				

R-4357 120324 12-03-24



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

FMC and Rhyme are
trademarks of FMC
Corporation® 2019