



BULLETIN
FIFRA Section 2(ee) Recommendation

DIMETRIC® EXT
EPA Reg. No. 1381-197
Tank Mix with
Gramoxone® SL 2.0, EPA Reg. No. 100-1431 or
Gramoxone® SL 3.0, EPA Reg. No. 100-1642
for Application to Alfalfa (Dormant Season)

FIFRA SECTION 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency.

This Bulletin expires 12/31/2026.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this Bulletin and the Dimetric EXT Herbicide EPA registered label. The EPA registered label must be in the possession of the user at the time of pesticide application.

Also refer to the Gramoxone SL 2.0 label or Gramoxone SL 3.0 label, whichever is applicable, for additional directions, weed species controlled, precautions and restrictions.

ALFALFA

Dimetric EXT plus Gramoxone® SL 2.0 (EPA Reg. No. 100-1431) Tank-mix
Dimetric EXT plus Gramoxone® SL 3.0 (EPA Reg. No. 100-1652) Tank-mix
For use in Colorado, Idaho, Montana, Nevada, Oregon, Utah, Washington, and Wyoming.

APPLICATION: Dimetric EXT plus Gramoxone SL 2.0 or Gramoxone SL 3.0 tank-mix application may be applied, during the dormant season, in aerial or ground spray equipment as a broadcast surface application to established (at least 1 year old) alfalfa for the control of certain grass and broadleaf weeds. Do not apply Dimetric EXT/Gramoxone SL 2.0 or Dimetric EXT/Gramoxone SL 3.0 tank-mix to regrowth (after grazing or cutting) that is more than 2 inches tall. Apply once per season. Do not apply following cuttings during growth season. Use a minimum of 10 gallons of water per acre with aerial spray equipment and a minimum of 20 gallons of water per acre with ground spray equipment. Add a nonionic spreader at label rates to the spray solution.

WEEDS CONTROLLED:

**Dimetric EXT (rates below) plus Gramoxone SL 2.0 (2 to 3 pt./acre) or
Dimetric EXT (rates below) plus Gramoxone SL 3.0 (1.3 to 2 pt./acre)**
tank-mix application will control established weeds.

1/3 to 1/2 Lb. of Dimetric EXT Per Acre			
Common Chickweed			
1/2 to 1 Lb. of Dimetric EXT Per Acre			
Bluegrass Cheat	Downy brome Field pennycress	Henbit Japanese brome	Rescuegrass Shepherdspurse
2/3 to 1 Lb. of Dimetric EXT Per Acre			
Blue mustard Common lambsquarters Flixweed Green foxtail Groundsel Jim Hill mustard	Kochia Little barley Marestail (Horseweed) Meadow salsify Pepperweed Prickly lettuce	Redroot pigweed Rough fleabane Ryegrass Smooth brome Sowthistle	Tansy mustard White cockle Wild oats Wild buckwheat Yellow rocket

Gramoxone SL 2.0 and Gramoxone SL 3.0 control weeds by contact activity.

	APPLICATION
DOSAGE/ACRE	
Dimetric EXT 1/3 to 1 Lb. Plus Gramoxone SL 2.0 2 to 3 pt.	Apply specified dosages of Dimetric EXT and Gramoxone SL 2.0 or Gramoxone SL 3.0 in at least 10 gallons of water per acre with aerial equipment or at least 20 gallons of water per acre with ground equipment. Do not apply this tank mix to alfalfa growth if more than 2 inches tall. For best weed control, apply when broadleaf weeds and grasses are 1-6 inches tall and are actively growing. Care should be taken to avoid overlaps. Do not apply more than 2/3 lb. of Dimetric EXT per acre on loamy sand soils. Reduced weed control may occur when extended dry conditions follow application of Dimetric EXT. Crop injury may occur if alfalfa is under stress conditions such as diseases, insect infestations, drought or winter injury or if Dimetric EXT is applied to alfalfa earlier than 12 months after seeding.
Dimetric EXT 1/3 to 1 Lb. Plus Gramoxone SL 3.0 1.3 to 2 pt.	

RESTRICTIONS:

Do not graze or harvest within 42 days after application.

In areas west of the Rockies, avoid the use of Dimetric EXT on soils with calcareous surface, soils with high levels of lime or sodium, and with a pH greater than 8.2.

Do not apply when weather conditions favor spray drift. Aerial application should not be made when wind speed is greater than 10 mph.

Do not use on sand soil.

Distributed by:

Winfield Solutions, LLC

P.O. Box 64589

St. Paul, MN 55164

12.13.2021

Dimetric is a registered trademark of Winfield Solutions, LLC

Gramoxone is a registered trademark of a Syngenta Group Company