

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



EPA Reg. No.: 279-3596

For early season disease protection in spring and winter wheat.

THIS 2(ee) RECOMMENDATION EXPIRES ON JUNE 30, 2023 AND MUST NOT BE USED AFTER THIS DATE

2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY

IN: ALABAMA, ARKANSAS, CONNECTICUT, DELAWARE, KENTUCKY, ILLINOIS, INDIANA, LOUISIANA, MAINE, MARYLAND, MICHIGAN, MINNESOTA, MISSISSIPPI, MONTANA, NEW HAMPSHIRE, NEW JERSEY, NORTH CAROLINA, NORTH DAKOTA, OHIO, PENNSYLVANIA, RHODE ISLAND, SOUTH CAROLINA, SOUTH DAKOTA, TENNESSEE, VERMONT, VIRGINIA, AND WEST VIRGINIA

Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING

CROP	PEST	RATE OF APPLICATION
Wheat (Spring and Winter)	Leaf, Stem and Stripe Rusts (<i>Puccinia</i> spp.) Powdery Mildew (<i>Blumeria graminis tritici</i>) Septoria Leaf and Glume Blotch (<i>Septoria</i> spp.) Tan Spot (<i>Pyrenophora tritici-repentis</i>)	3 - 4 fl. oz./A

DIRECTIONS FOR USE:

Apply Topguard® EQ fungicide in early spring from tillering to jointing growth stages (Feekes stages 3-7) to provide protection against listed early season leaf diseases. Topguard EQ fungicide may be tank-mixed with adjuvants, fertilizer, insecticides (such as **Mustang® MAXX** insecticide) or herbicides (such as **Harmony® Extra SG** herbicide with TotalSol® soluble granules or **Finesse®** Cereal and Fallow herbicide) to implement disease, insect and weed management. An application of Topguard EQ fungicide may be required at flag leaf (Feekes 8-9) for control of foliar diseases as specified by product label. Apply in a minimum of 2 gallons per acre for aerial application and 10 gallons per acre by ground application.

RESTRICTIONS:

- Do not apply more than 2 applications per year.
- Do not apply more than 0.228 lb. ai of flutriafol/A per year.
- Do not apply more than 0.4 lb. ai of azoxystrobin per acre per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Intervals (PHI): Do not apply within 30 days of harvest when grown for grain, 15 days of harvest when grown for hay, 7 days of harvest when grown for forage.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED TOPGUARD EQ FUNGICIDE, EPA REG.NO. 279-3596, LABEL MUST BE FOLLOWED. THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

R-4218 040919

FMC, Topguard, Harmony, TotalSol and Mustang
are trademarks of FMC Corporation
©2018 FMC Corporation. All rights reserved