

**THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.**



**EPA Reg. No.: 279-3596**

## 2(ee) Recommendation

### FOR DISTRIBUTION AND USE ONLY IN:

ALABAMA, ARKANSAS, COLORADO, CONNECTICUT, DELAWARE, GEORGIA, IDAHO, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, LOUISIANA, MAINE, MARYLAND, MASSACHUSETTS, MICHIGAN, MINNESOTA, MISSISSIPPI, MISSOURI, MONTANA, NEBRASKA, NEW HAMPSHIRE, NEW JERSEY, NEW YORK, NORTH CAROLINA, NORTH DAKOTA, OHIO, OKLAHOMA, OREGON, PENNSYLVANIA, RHODE ISLAND, SOUTH CAROLINA, SOUTH DAKOTA, TENNESSEE, VERMONT, VIRGINIA, WEST VIRGINIA, WASHINGTON, WISCONSIN AND WYOMING

Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING**

### For Fall and Spring Disease Management in Spring and Winter Wheat and Barley

*THIS 2(ee) RECOMMENDATION EXPIRES ON DECEMBER 31, 2028 AND MUST NOT BE USED AFTER THIS DATE*

CROP	PEST	RATE OF APPLICATION
Wheat (Spring and Winter) Barley	Leaf, Stem and Stripe Rusts ( <i>Puccinia</i> spp.) Powdery Mildew ( <i>Blumeria graminis tritici</i> ) Septoria Leaf and Glume Blotch ( <i>Septoria</i> spp.) Tan Spot ( <i>Pyrenophora tritici-repentis</i> )	3 - 4 fl. oz./A

#### DIRECTIONS FOR USE:

- Apply Topguard® EQ fungicide in the fall after emergence and/or in early spring from tillering to jointing growth stages (Feekes stages 2-7) to provide protection against listed early season leaf diseases.
- Topguard EQ fungicide may be tank-mixed with adjuvants, fertilizer, labeled insecticides (such as **Mustang® MAXX** insecticide) or herbicides (such as **Affinity®** brand herbicides with TotalSol® soluble granules, **Harmony® Extra SG** herbicide with TotalSol soluble granules or **Finesse®** Cereal and Fallow herbicide) to implement disease, insect and weed management.
- An application of Topguard EQ fungicide may be required at flag leaf (Feekes 8 - 9) for control of foliar diseases as specified by product label.
- Apply in a minimum of 2 gallons per acre for aerial application and 10 gallons per acre by ground application.

#### RESTRICTIONS:

- Do not apply more than 2 applications per year.
- Do not apply more than 0.228 lb. ai of flutriafol/A per year.
- Do not apply more than 0.4 lb. ai of azoxystrobin per acre per year.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Intervals (PHI): Do not apply within 30 days of harvest when grown for grain, 15 days of harvest when grown for hay, 7 days of harvest when grown for forage.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED TOPGUARD EQ FUNGICIDE, EPA REG.NO. 279-3596, LABEL MUST BE FOLLOWED. THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.



FMC Corporation  
2929 Walnut Street  
Philadelphia, PA 19104

R-4831 070723  
FMC, Topguard, Harmony, TotalSol, Mustang  
Finesse and Affinity are trademarks of FMC Corporation  
©2023 FMC Corporation. All rights reserved