



## FIFRA 2(ee) Recommendation

EPA Reg. No. 279-3594

**FOR DISTRIBUTION AND USE  
ONLY IN THE FOLLOWING  
STATES:TX**

**THIS RECOMMENDATION, WHICH  
CONTAINS ADDITIONAL DIRECTIONS  
FOR USE, IS MADE AS PERMITTED  
UNDER FIFRA SECTION 2(ee) AND HAS  
NOT BEEN SUBMITTED TO OR  
APPROVED BY THE US EPA.**

This recommendation for Topguard Terra Fungicide is valid until December 31, 2025, or until withdrawn, canceled or suspended.

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.**

**THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**

**For Use in Cotton as Preplant injection, Preemergence, Early Postemergence  
and Post Directed Application for Management of Cotton Root Rot  
(*Phymatotrichopsis omnivora*).**

### Directions For Use

#### Method of Application

- Topguard® Terra fungicide may be injected into row up to 30 days prior to planting. Fungicide should be placed 2-3 inches directly below seed placement zone.
- Topguard® Terra fungicide may be applied in 3 – 4 inch band on top of row prior to cotton emergence.
- Topguard® Terra fungicide may be applied to emerged cotton either as a 4 inch band over-the-top of cotton not larger than 2 true leaves.
- Topguard Terra may be applied as a directed spray which should be concentrated in a band that is no wider than 4 inches and directed at the base of the plant. Cotton should be no larger than 4 true leaves.



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104  
215-299-6000

## Rates and Timing

Application Timing	Use Rate (fl oz/A (lb ai/A))	Comments
<ul style="list-style-type: none"> <li>➤ <b>Preplant Injected</b> into row up to 30 days prior to planting.</li> <li>➤ <b>Preemergence</b> in 3-4 inch band behind the planter press wheel.</li> <li>➤ <b>Postemergence (Over the Top)</b> of cotyledon to 2 leaf cotton.</li> <li>➤ <b>Directed-postemergence</b> 2-4 leaf cotton.</li> </ul>	4-8 fl oz/A (0.13 – 0.26 lb ai/A)	Use higher rates on irrigated fields or where high infestations of cotton root rot are common. Rainfall or irrigation is required to move Topguard Terra into infection zone.

R-4557 030221



FMC Corporation  
 2929 Walnut St.  
 Philadelphia, PA 19104  
 215-299-6000