

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA



EPA Reg. No. 279-3557

2(ee) Recommendation

**FOR DISTRIBUTION AND USE ONLY IN
Colorado, Kansas, Missouri, Montana, Nebraska,
North Dakota, Oklahoma, and South Dakota**

**THIS PRODUCT BULLETIN CONTAINS NEW OR
ADDITIONAL DIRECTIONS FOR USE WHICH ARE
RECOMMENDED BY FMC CORPORATION UNDER
FIFRA SECTION 2(ee)**

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS
PRODUCT IN A MANNER INCONSISTENT WITH IT
LABELING**

For early season disease management in winter wheat.

Crop	Pest	Rate of Application
Winter Wheat	Common and stripe rusts (<i>puccinia spp.</i>) Powdery mildew (<i>blumeria graminis tritici</i>) Leaf blotch (<i>septoria spp.</i>)	5 – 7 fl. oz./A

Application Instructions and Restrictions

- Apply in early spring tillering to early joint (feekes stages 3-7).
- Do not apply more than 0.228 lb. ai of flutriafol/A per year.
- Do not apply more than 2 applications per year.
- Do not apply more than 28 fl. oz. of product/A per year.

Follow with a flag leaf application (feekes 9-10) of **Topguard EQ** or **Preemptor™ SC** fungicides per the product label of for full season control of foliar diseases.

Topguard fungicide may be mixed with adjuvants, insecticides (such as **Mustang® Maxx** insecticide) or herbicides (such as **Finesse®** Cereal and Fallow herbicide) to provide insect or weed control and disease management in one application.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED TOPGUARD FUNGICIDE LABEL (EPA REG. NO.279-3557) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104
DOC 4-16-2018

FMC, Topguard, Preemptor,
Mustang and Finesse are
trademarks of FMC Corporation
Copyright © 2018