

RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

BIFENTHRIN	GROUP	3A	INSECTICIDE
B. AMYLOLIQUEFACIENS	GROUP	44	FUNGICIDE

Supplemental Labeling

This supplemental label expires on December 31, 2024 and must not be used or distributed after this date.



For At-Plant use on Sunflower

Liquid insecticide/Fungicide for use in a 3RIVE 3D system only. Do not apply by air.

EPA Reg. No. 279-3632

Active Ingredient:	By Wt.
Bifenthrin: *	15.67%
Bacillus amyloliquefaciens strain D747**	5.50%
Other Ingredients:	78.83%
.....	100.0%

*Cis isomers 97% minimum, trans isomers 3% maximum

** Contains a minimum of 1.0×10^{10} colony-forming units (cfu) per milliliter of product

This product contains 1.4 lbs. bifenthrin per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER
INCONSISTENT WITH ITS LABELING.**

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA
REGISTERED LABEL ARE TO BE FOLLOWED.**

This Supplemental labeling must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for Ethos 3D Insecticide/Fungicide before applying. Use of Ethos 3D Insecticide/Fungicide according to this supplemental labeling is subject to the use precautions and limitations imposed by the label affixed to the container of Ethos 3D Insecticide/Fungicide. Carefully follow all precautionary statements and application use directions.



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

3RIVE 3D System Information: Ethos 3D Insecticide/Fungicide must be applied with a 3RIVE 3D system. Refer to the 3RIVE 3D equipment and user manual for information on system operation and calibration.

SUNFLOWER (Crop Subgroup 20B)

Calendula, Castor Oil Plant, Chinese Tallowtree, Euphorbia, Evening Primrose, Jojoba, Niger Seed, Rose Hip, Safflower, Stokes Aster, Tallowwood, Tea Oil Plant, Vernonia, cultivars, varieties, and/or hybrids of these

At-Plant

PEST/DISEASE	USE RATES		DIRECTIONS
	Fluid oz/1000 Linear ft. ¹	Pound bifenthrin/acre	
Wireworm Grape colaspis White grub species Seedcorn maggot Root aphids Army cutworm Cutworm species	0.21 – 1.05	0.04 - 0.2	Apply in-furrow with the seed. Ethos 3D Insecticide/Fungicide must be applied with a 3Rive 3D system.
² Suppression of "Damping off," seedling blights, and root or crown diseases caused by <i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i> , or <i>Phytophthora</i>			
At-Plant Restrictions:			
<ul style="list-style-type: none"> Do not apply more than 0.2 lb ai/A (16.0 fl oz/A) per year as an at-plant application. Do not make more than 1 application per year. 			

¹ Based on 30" row spacing

² Use highest labeled rate for disease management; however, under moderate to severe disease pressure, or when environmental conditions and plant stage are conducive to rapid disease development, mix Ethos 3D Insecticide/Fungicide with other fungicides for improved performance.

FMC, ETHOS and 3RIVE 3D are trademarks of FMC Corporation.
© 2021 FMC Corporation. All rights reserved.

R-4631 012522 11-19-2021