



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-3603

FOR DISTRIBUTION AND USE ONLY IN: Arkansas, Connecticut, Delaware, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nebraska, New Hampshire, New Jersey, North Dakota, Ohio, Oklahoma, Pennsylvania, Rhode Island, Utah, South Dakota, Tennessee, Vermont, Virginia, West Virginia, Wisconsin and Wyoming

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation for Lucento™ Fungicide is valid until May 25, 2025, or until withdrawn, canceled or suspended.

Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

For Tar Spot Management In Corn (Field, Corn Grown For Seed And Popcorn)

Directions For Use

Apply at the onset of disease up through the R4 growth stage. Repeat applications at 7-14 day intervals if needed. Use the higher use rate and shorter interval when disease pressure is high and conditions are favorable for disease development.



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000

CROP	PEST	RATE OF APPLICATION
Corn (Field, Corn Grown for Seed and Popcorn)	Tar Spot (<i>Phyllachora maydis</i>)	5.0 - 5.5 fl. oz./A
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • Do not apply after the R4 growth stage (early dough) of corn. • Do not exceed 5.5 fl. oz./A in a single application. • Maximum 2 applications per year. • Do not exceed 11 fl. oz. of product/A per year. • Do not apply more than 0.228 lb. ai of flutriafol/A per year from all flutriafol containing products. • Do not apply more than 0.132 lb. ai of bixafen/A per year from all bixafen containing products. • Do not use an adjuvant after the V8 stage and prior to the VT stage of corn. An adjuvant may be used at any other growth stage. <p>Restricted Entry Interval (REI): The REI for detasselling field corn and popcorn grown for seed is 5 days. The REI for all other activities is 12 hours.</p> <p>Preharvest Interval: May be applied up to 10 days before harvest for forage, and 30 days before harvest for grain or stover.</p>		

R-4443 041620



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000