

# FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9615

# FOR DISTRIBUTION AND USE ONLY IN THE STATE OF NEW YORK

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation for EXIREL® insect control is valid until May 31, 2025, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

# FOR CONTROL OF APPLE LEAF CURLING MIDGE IN APPLES IN THE STATE OF NEW YORK

## **Directions For Use**

# Application Rates, Timing, Method and Pest Controlled

EXIREL® insect control is recommended for control of apple leaf curling midge (Dasineura mali).

Apply EXIREL insect control at 10 - 20.5 fl. oz./A (0.065 - 0.133 lb ai/A) as a foliar spray.

Initiate foliar applications at the first sign of egg laying or at early egg hatch. Two sequential applications, 10 to 14 days apart, may be needed in areas of high pest pressure. Follow local university recommendations for treatment thresholds and additional instructions on optimum application timing.

Thorough spray coverage is essential for best performance. Select an appropriate spray volume based on the size of the trees and foliage density. Do not apply less than 30 gallons of water per acre by ground.

See "Tank Mixtures and Crop Safety" section on the EXIREL insect control label for more information, and for guidance regarding crop response for tested tank mixtures.



### **SPRAY PREPARATION**

**Acidification of spray tank:** If the pH of the spray tank after all products have been added and mixed is above pH 8, adjust to pH 8 or less using a registered acidifying agent. If the spray tank pH is 8 or less, no adjustment is necessary. Spray tanks of pH 8 or less can be held up to 8 hours before spraying. Do not store the spray mixture overnight in the spray tank.

### **RESTRICTIONS AND PRECAUTIONS**

- Minimum interval between treatments is 7 days.
- Do not apply a total of more than 0.4 lb ai/A Cyazypyr® active or cyantraniliprole containing products per calendar year.
- Make no more than 3 applications of Exirel insect control or other Group 28 insecticide within a single generation of the target pest on a crop.
- The preharvest interval (PHI) is 3 days.
- The field re-entry interval (REI) is 12 hours.

### IMPORTANT NOTICE

BEFORE BUYING OR USING FMC PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY, AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of Federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This Bulletin contains supplemental instructions for use of the referenced FMC product. Follow these instructions carefully. This Bulletin must be in the possession of the User at the time of application. Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using the FMC product(s) covered by this Bulletin. FMC will not be responsible for losses or damages resulting from use of this product in any manner not specifically stated by FMC on the Section 3 label, or other labels, or this Bulletin. User assumes all risks associated with any such nonspecified use. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability of the Section 3 Federal product label.

R-4399 033120 03-30-20

FMC, the FMC logo, EXIREL and CYAZYPYR are Trademarks of FMC Corporation or an affiliate.  $@2020\,\mathrm{FMC}$  Corporation

