



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9615

**FOR DISTRIBUTION AND USE
ONLY IN THE STATE OF FLORIDA**

**THIS RECOMMENDATION, WHICH CONTAINS
ADDITIONAL DIRECTIONS FOR USE, IS
MADE AS PERMITTED UNDER FIFRA
SECTION 2(ee) AND HAS NOT BEEN
SUBMITTED TO OR APPROVED BY THE US
EPA.**

This recommendation for EXIREL® insect control is valid until March 1, 2026, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

CONCENTRATION BASED RATES FOR CONTROL OF ASIAN CITRUS PSYLLID, CITRUS LEAFMINER, AND DIAPREPES ROOT WEEVIL ADULTS IN YOUNG RESET OR SOLID-SET CITRUS TREES IN THE STATE OF FLORIDA

Directions for Use

EXIREL insect control is recommended for control of Asian citrus psyllid, citrus leafminer and Diaprepes root weevil adults in young reset or solid-set citrus trees (crop group 10-10) up to 24 months old.

Application Rate: Apply EXIREL insect control at 16-20.5 fluid ounces per 100 gallons of water. For best performance, use of an adjuvant such as horticultural spray oil at 1 gallon per 100 gallons of spray volume (1% v/v) is recommended. Do not apply more than 0.133 lb. of CYAZYPYR® active per acre per application (20.5 fl. oz./A EXIREL insect control) or a total of more than 0.4 lb of CYAZYPYR active or other cyantraniliprole containing products per acre per calendar year.

Application Timing: EXIREL insect control should be timed to protect new flush. Applications should be made early during the feather flush stage when the new flush is no more than 40% expanded and before heavy insect damage is present.

Spray Volume: Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of trees and density of foliage to ensure adequate coverage to wet the foliage. The amount of treatment solution per acre will vary based on the size of trees being treated and planting density. When using a rate of 20.5 fl. oz. EXIREL insect control per 100 gallons of water, do not apply more than 100 gallons of spray solution per acre.



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104

Application Equipment: Applications can be made with a handgun or hoop boom sprayer to young reset or solid-set citrus trees in sufficient spray volume to wet all plant foliage.

SPRAY PREPARATION

Acidification of the spray tank: If the pH of the spray tank after all products have been added and mixed is above pH 8, adjust to pH 8 or less using a registered acidifying agent. If the spray tank pH is 8 or less no adjustment is necessary. Spray tanks of pH 8 or less can be held up to 8 hours before spraying. Do not store the spray mixture overnight in the spray tank.

See the RESISTANCE MANAGEMENT section of the EXIREL insect control Federal label for additional guidance regarding insecticide resistance management.

See 'Tank Mixtures and Crop Safety' section on the EXIREL insect control label for more information and for guidance regarding crop response for tested tank mixtures.

RESTRICTIONS

Do not apply EXIREL insect control to the soil or through drip irrigation systems.

Do not apply a total of more than 0.4 lb a.i./A. of CYAZYPYR active or cyantraniliprole-containing products per calendar year whether applications are foliar or soil-applied.

The minimum retreatment interval is 7 days.

The PHI (pre-harvest interval) is 1 day.

The REI (re-entry interval) is 12 hours.

R-4572 022421 05-25-21

FMC, the FMC logo, CYAZYPYR and EXIREL are Trademarks of FMC Corporation or an affiliate.
©2021 FMC Corporation



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104