

SUPPLEMENTAL LABELING



EXIREL[®]
INSECT CONTROL

CYANTRANILIPROLE	GROUP	28	INSECTICIDE
------------------	-------	-----------	-------------

WITH CYAZYPYR[®] active

EPA Reg. No. 279-9615

This label expires on September 29, 2027

FOR FOLIAR APPLICATIONS TO HOPS FOR PEST MANAGEMENT OF HOP FLEA BEETLE (CHEWING INSECT), AIDING IN OPTIMIZATION OF THE CROP'S POTENTIAL.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT

BEFORE USING EXIREL[®] INSECT CONTROL, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This supplemental label contains new or additional instructions for use of this product which does not appear on the package label. Follow the instructions carefully.

This label must be in the possession of the user at the time of pesticide application.

GENERAL INFORMATION

EXIREL[®] insect control is a suspoemulsion (oil in water emulsion) that can be applied as a foliar spray on labeled crops to control listed insects.

EXIREL[®] insect control is specially formulated for maximum performance by foliar applications to hops for pest management of hop flea beetle (chewing insect), aiding in optimization of the crop's potential.

Sold By



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

Directions for Use

Crop	Application Method	Target Pest	EXIREL® insect control RATE		PHI (pre-harvest interval) (days)	REI (re-entry interval) (hours)
			Lb. ai per acre	fluid ounces product per acre		
Hops*	Foliar	Hop flea beetle	0.088 - 0.133	13.5 - 20.5	1	12
<p>*Not registered for use by CA. Minimum application interval between treatments is 5 days. Do not make more than 3 applications per calendar year. Do not apply a total of more than 0.4 lb ai/A of CYAZYPYR® active or cyantraniliprole containing products per calendar year whether applications are made to the soil or foliarly. Spray Volume: Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of trees or plants and density of foliage. Apply in a least 30 gallons per acre by ground application equipment. For best performance use an effective adjuvant. See "Use of Adjuvants" section. When using EXIREL® insect control in tank mixtures, it is recommended that a small area be tested to demonstrate safety before using in large areas. See "Tank Mixtures and Crop Safety" section for more information.</p>						

FMC, the FMC logo, Exirel and Cyazypyr are trademarks of FMC Corporation and/or an affiliate.
 © 2024 FMC Corporation. All rights reserved.

(Derived from Exirel insect control label SL-4712A 103124 08-29-24)

R-4713A 032326 08-29-24