

SUPPLEMENTAL LABELING



EXIREL®



INSECT CONTROL

CYANTRANILIPROLE	GROUP	28	INSECTICIDE
------------------	-------	----	-------------

WITH CYAZYPYR® active

EPA Reg. No. 279-9615

This label expires on September 29, 2027

FOR FOLIAR APPLICATIONS TO PAPAYA, HERBS (FRESH AND DRIED), SPICES, AND GREENHOUSE GROWN LETTUCE FOR PEST MANAGEMENT OF SUCKING AND CHEWING INSECTS THAT CAN VECTOR CERTAIN PLANT DISEASES, AIDING IN OPTIMIZATION OF THE CROP'S POTENTIAL.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT

BEFORE USING EXIREL® INSECT CONTROL, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This supplemental label contains new or additional instructions for use of this product which does not appear on the package label. Follow the instructions carefully.

This label must be in the possession of the user at the time of pesticide application.

GENERAL INFORMATION

EXIREL® insect control is a suspoemulsion (oil in water emulsion) that can be applied as a foliar spray on labeled crops to control listed insects. EXIREL® insect control is specially formulated for maximum performance by foliar applications to basil, dill, mint and papaya for pest management of sucking and chewing insects that can vector certain plant diseases, aiding in optimization of the crop's potential.

Sold By



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

Directions for Use

Crop	Application Method	Target Pest	EXIREL® insect control RATE		PHI (pre-harvest interval) (days)	REI (re-entry interval) (hours)
			Lb. ai per acre	fluid ounces product per acre		
Papaya	Foliar	Thrips (foliage feeding only)§	0.088 - 0.133	13.5 - 20.5	1	12
<p>Minimum application interval between treatments is 14 days. Do not make more than 3 applications per calendar year. Do not apply a total of more than 0.4 lb ai/A of CYAZYPYR® active or cyantraniliprole containing products per calendar year whether applications are made to the soil or foliarly. §-Suppression only. For best results, use the highest rate listed. Use as part of an effective thrips control program. Rotate with products with different modes of action. Spray Volume: Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of trees or plants and density of foliage. Apply in a least 30 gallons per acre by ground application equipment. For best performance use an effective adjuvant. See "Use of Adjuvants" section. The crop safety of EXIREL® insect control in tank mixture has not been evaluated on this crop or crop group. When using EXIREL® insect control in tank mixtures, it is recommended that a small area be tested to demonstrate safety before using in large areas. See "Tank Mixtures and Crop Safety" section for more information.</p>						

Crop	Application Method	Target Pest	EXIREL® insect control RATE		PHI (pre-harvest interval) (days)	REI (re-entry interval) (hours)
			Lb. ai per acre	fluid ounces product per acre		
Herbs, fresh leaves (Crop Subgroup 25A) including Curry leaf	Foliar	Asian citrus psyllid	0.088 - 0.133	13.5 - 20.5	1	12
	<p>Minimum application interval between treatments is 5 days. Do not make more than 3 applications per calendar year. Do not apply a total of more than 0.4 lb ai/A of CYAZYPYR® active or cyantraniliprole containing products per calendar year whether applications are made to the soil or foliarly. Spray Volume: Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of trees or plants and density of foliage. Apply in a least 20 gallons per acre by ground application equipment. When using EXIREL® insect control in tank mixtures, it is recommended that a small area be tested to demonstrate safety before using in large areas. See "Tank Mixtures and Crop Safety" section for more information.</p>					

Crop	Application Method	Target Pest	EXIREL® insect control RATE		PHI (pre-harvest interval) (days)	REI (re-entry interval) (hours)
			Lb. ai per acre	fluid ounces product per acre		
Herbs, dried leaves (Crop Subgroup 25B) including Curry leaf	Foliar	Asian citrus psyllid	0.088 - 0.133	13.5 - 20.5	1	12
	<p>Minimum application interval between treatments is 5 days. Do not make more than 3 applications per calendar year. Do not apply a total of more than 0.4 lb ai/A of CYAZYPYR® active or cyantraniliprole containing products per calendar year whether applications are made to the soil or foliarly. Spray Volume: Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of trees or plants and density of foliage. Apply in a least 20 gallons per acre by ground application equipment. When using EXIREL® insect control in tank mixtures, it is recommended that a small area be tested to demonstrate safety before using in large areas. See "Tank Mixtures and Crop Safety" section for more information.</p>					

Crop	Application Method	Target Pest	EXIREL® insect control RATE		PHI (pre-harvest interval) (days)	REI (re-entry interval) (hours)
			Lb. ai per acre	fluid ounces product per acre		
Spices (Crop Group 26) including Kaffir lime	Foliar	Asian citrus psyllid	0.088 - 0.133	13.5 - 20.5	1	12
	<p>Minimum application interval between treatments is 5 days. Do not make more than 3 applications per calendar year. Do not apply a total of more than 0.4 lb ai/A of CYAZYPYR® active or cyantraniliprole containing products per calendar year whether applications are made to the soil or foliarly. Spray Volume: Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of trees or plants and density of foliage. Apply in a least 20 gallons per acre by ground application equipment. When using EXIREL® insect control in tank mixtures, it is recommended that a small area be tested to demonstrate safety before using in large areas. See "Tank Mixtures and Crop Safety" section for more information.</p>					

Crop	Application Method	Target Pest	EXIREL® insect control RATE		PHI (pre-harvest interval) (days)	REI (re-entry interval) (hours)
			Lb. ai per acre	fluid ounces product per acre		
Greenhouse Grown Lettuce	Foliar*	Beet armyworm Corn earworm Fall armyworm Western yellowstriped armyworm	0.045 - 0.088	7 - 13.5	1	12
		Cabbage looper	0.065 - 0.11	10 - 17		
		Cabbage aphid False cabbage aphid Green peach aphid Thrips (foliage feeding only)§ Turnip aphids Whitefly	0.088 - 0.133	13.5 - 20.5		
<p>Minimum application interval between treatments is 5 days. Do not make more than 3 applications per calendar year. Do not apply a total of more than 0.4 lb ai/A of CYAZYPYR® active or cyantraniliprole containing products per calendar year whether applications are made to the soil or foliarly. Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of plants and density of foliage. Use the higher rate on large plants or dense foliage *- For best performance, use with an effective adjuvant. See "Use of Adjuvants" section. § - Suppression only. Use as part of an effective thrips control program. Rotate with products with different modes of action. Begin making applications to thrips when populations are low. If populations are above threshold, use an effective thrips knockdown product before applying EXIREL® insect control. Precautions when using EXIREL® insect control in tank mixes in lettuce: Tank mixes of EXIREL® insect control with Alette® fungicide (fosetyl-al) + oil adjuvant may result in adverse crop response. When using EXIREL® insect control in tank mixtures, it is recommended that a small area be tested to demonstrate safety before using in large areas. See "Tank Mixtures and Crop Safety" section for more information.</p>						

FMC, the FMC logo, Exirel and Cyazypyr are trademarks of FMC Corporation and/or an affiliate.
© 2024 FMC Corporation. All rights reserved.

(Derived from Exirel insect control label SL-4712A 103124 08-29-24)

R-4713A 071825 08-29-24