

FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9587

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF OREGON AND WASHINGTON

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation for AVAUNT® insecticide is valid until May 31, 2026, or until withdrawn, canceled, or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR CONTROL OF BLACK VINE WEEVIL (ADULT) IN CRANBERRY

Directions For Use

AVAUNT insecticide is recommended as a foliar application to control adult black vine weevil in cranberry.

Apply AVAUNT insecticide at 6.0 oz/acre (0.11 lb a.i./acre) as a foliar spray based on locally established treatment thresholds.

Black vine weevil adults are nocturnal feeders - it is important to monitor adult emergence by regular sweeping or trapping in the evening hours. Make repeat applications on a 7 to 10 day schedule if monitoring indicates continued adult feeding activity. Broadcast applications may need supplemental spot treatments in localized areas of heavy insect pressure. Allow 5 to 7 days to achieve maximum efficacy.

RESTRICTIONS:

Do not apply more than 24 oz AVAUNT insecticide (0.44 lb a.i.) per acre per crop.

Make no more than 3 applications per acre per season.

The minimum interval between sprays is 7 days.

The preharvest interval is 30 days.

Do not apply to flow through bogs or allow release of irrigation water from bogs for at least 1 day following application.



R-4579 060421 05-25-21

FMC, the FMC logo and Avaunt are trademarks of FMC Corporation or an affiliate. $@2021\ \text{FMC}$ Corporation

