FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9596



FOR DISTRIBUTION AND USE ONLY IN THE STATE OF MASSACHUSETTS

THIS RECOMMENDATION, WHICH CONTAINS
ADDITIONAL DIRECTIONS FOR USE, IS MADE AS
PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS
NOT BEEN SUBMITTED TO OR APPROVED BY THE US
EPA.

This recommendation for STEWARD® EC insecticide is valid until December 31, 2025, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR CONTROL OF TOBACCO FLEA BEETLE IN TOBACCO

CROP	APPLICATION METHOD	TARGET PEST	RATE	PHI (Days)	REI (Hours)
Tobacco	Foliar	Tobacco Flea Beetle (<i>Epitrix hiritipennis</i>)	9.2 – 11.3 fl oz/A (0.09 - 0.11 lb ai/A)	14	12

Directions For Use:

- Steward EC insecticide is recommended as a foliar application for control of tobacco flea beetle.
- Higher application volume (GPA) will provide better coverage and performance. Select a spray volume appropriate for the size of plants and density of foliage being treated. A minimum of 10 GPA by ground application equipment is recommended.
- Use high rate (11.3 fl oz/A) for optimal control and when heavier infestations are present.
- See the Resistance Management section of the Steward EC insecticide Section 3 Federal label for additional guidance regarding insecticide resistance management.



RESTRICTIONS AND PRECAUTIONS

- Do not apply more than 45 fl oz/A Steward EC insecticide or 0.44 lb ai/A of indoxacarb-containing products per year.
- Do not make more than 4 applications per acre per crop.
- The minimum interval between treatments is 5 days.

FMC and Steward are trademarks of FMC Corporation or an affiliate. ©2020 FMC Corporation

R-4550 020521 01-29-21