



INDOXACARB	GROUP	22A	INSECTICIDE
------------	-------	-----	-------------

EPA Reg. No. 279-9596

SUPPLEMENTAL LABELING

**FOR FOLIAR APPLICATIONS TO TOBACCO TO CONTROL
TOBACCO BUDWORM, TOBACCO HORNWORM, TOMATO
HORNWORM AND TOBACCO SPLITWORM**

**THIS SUPPLEMENTAL LABEL EXPIRES 11/30/2023 AND MUST NOT BE USED OR
DISTRIBUTED AFTER THIS DATE**

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read the label affixed to the container for Steward EC insecticide before applying.

Use of Steward EC insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Steward EC insecticide.

IMPORTANT

**BEFORE USING STEWARD EC insecticide, READ AND FOLLOW ALL APPLICABLE DIRECTIONS;
RESTRICTIONS; AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which does not appear on the package label. Follow the instructions carefully.

This label must be in the possession of the user at the time of pesticide application.

Sold By
FMC
FMC Corporation 2929
Walnut Street
Philadelphia, PA 19104

STEWARD EC insecticide Rate Per Acre

Crops	Insects	lb ai	fl oz	Acres Treated per gal of STEWARD EC insecticide	Last Application (Days to Harvest)	REI
Tobacco	Tobacco budworm Tobacco hornworm Tomato hornworm Tobacco splitworm	0.065 - 0.11	6.7 - 11.3	11.5 - 19	14	12 hrs.

TOBACCO RESTRICTIONS

The minimum interval between treatments is 5 days.

Make no more than 4 applications per acre per crop.

Do not apply more than 45 fl oz/A of STEWARD EC insecticide or 0.44 lb ai/A of indoxacarb-containing products per calendar year.

Note: Use intermediate to high rates within the listed rate range (9.2 to 11.3 fl oz/A) on heavier infestations or when later instar larvae are present.

(Derived from STEWARD EC Insecticide Label SL-4292 071720 11-23-20)

R-4298 111920 11-23-20