THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA



FIFRA 2(ee) Recommendation EPA Reg. No. 279-9596

FOR DISTRIBUTION AND USE ONLY IN: Colorado, Kansas, Illinois, Indiana, Missouri, Nebraska, North Dakota, Oklahoma, and South Dakota

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

This recommendation for STEWARD EC insecticide is valid until 12/31/2029, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR USE OF STEWARD EC INSECTICIDE FOR CORN LEAFHOPPER ON CORN

CROP	PEST	RATE OF APPLICATION
Corn (field) Corn (grown for seed) Corn (pop)	Corn Leafhopper (<i>Dalbulus maidis</i>)	6.0 – 11.3 fl. oz./A
APPLICATION INSTRUCTIONS AND RESTRICTIONS:		
The minimum interval between treatments is 5 days.		
 Make no more than 2 applications per acre per crop. 		
Do not apply more than 22.6 fl oz/A of STEWARD EC insecticide or 0.22 lb ai/A of indoxacarb-		
containing products per calendar year.		
 Apply in a minimum of 3 GPA (aerial) and 10 GPA (ground) for optimal coverage and efficacy. 		
Application by chemigation systems is allowed.		
The REI is 12 hours.		
 Interval between last application and harvest is: 1 day for forage, fodder, and silage and 14 days 		
for grain and stover.		
STEWARD EC insecticide may be mixed with adjuvants, fungicides, or herbicides to provide disease or		
weed control and insect management in one application.		

R-4928 111224 11-12-24

STEWARD and FMC – Trademarks of FMC or an affiliate ©2024 FMC Corporation. All rights reserved.

