



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9616

**FOR DISTRIBUTION AND USE
ONLY IN THE STATES OF
GEORGIA, KENTUCKY, NORTH
CAROLINA, PENNSYLVANIA,
SOUTH CAROLINA, TENNESSEE
AND VIRGINIA**

**THIS RECOMMENDATION, WHICH CONTAINS
ADDITIONAL DIRECTIONS FOR USE, IS MADE
AS PERMITTED UNDER FIFRA SECTION 2(ee)
AND HAS NOT BEEN SUBMITTED TO OR
APPROVED BY THE US EPA.**

This recommendation for VERIMARK® insect control is valid until May 31, 2026, or until withdrawn, canceled, or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR THE SUPPRESSION OF TOMATO SPOTTED WILT VIRUS INCIDENCE IN TOBACCO VECTORED BY THRIPS

Directions for Use

VERIMARK insect control used in programs with imidacloprid is recommended for the suppression of tomato spotted wilt virus in tobacco when vectored by thrips.

Apply VERIMARK insect control at 13.5 fluid ounces per acre as a transplant tray drench or transplant water application. An application of a registered product that contains imidacloprid as a tray drench or transplant water at labeled rates, typically 0.028 - 0.04 lb ai/1000 plants, should also be made. VERIMARK insect control and imidacloprid can be applied together using the same application method or separately using different application methods. When applying VERIMARK insect control as a tray drench, apply no earlier than 72 hours prior to planting in the field. Read and follow label directions for transplant tray drench and transplant water application methods as described on the section 3 label of VERIMARK insect control and the imidacloprid containing product.

VERIMARK insect control impairs insect muscle function, resulting in rapid cessation of feeding and limiting the ability of insects to move, which can reduce transmission and spreading of insect vectored diseases.



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104

See the RESISTANCE MANAGEMENT section of the VERIMARK insect control Federal label for additional guidance regarding insecticide resistance management.

See 'Tank Mixtures and Crop Safety' section on the VERIMARK insect control label for more information and for guidance regarding crop response for tested tank mixtures.

USE RESTRICTIONS

Do not apply more than 13.5 fl oz (0.176 lb ai/A) of VERIMARK insect control at planting.

Do not apply a total of more than 0.4 lb ai/A of CYAZYPYR® active or cyantraniliprole containing products per acre per calendar year whether the applications are made to the soil or foliarly.

Re-entry interval after a Verimark insect control application is 4 hours.

R-4575 031021 05-25-21

FMC, the FMC logo, CYAZYPYR and VERIMARK are Trademarks of FMC Corporation or an affiliate.

©2021 FMC Corporation



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104