



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9616

FOR DISTRIBUTION AND USE
ONLY IN THE STATE OF NEW YORK

THIS RECOMMENDATION, WHICH
CONTAINS ADDITIONAL DIRECTIONS
FOR USE, IS MADE AS PERMITTED
UNDER FIFRA SECTION 2(ee) AND HAS
NOT BEEN SUBMITTED TO OR
APPROVED BY THE US EPA.

This recommendation for VERIMARK® Insect Control is valid until December 31, 2030, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR SUPPRESSION OF ONION MAGGOT AND SEEDCORN MAGGOT IN ONIONS (CROP GROUP 3-07)

Directions For Use

APPLICATION METHOD, RATES AND TIMINGS

Tray Drench

Apply VERIMARK insect control at 13.5 fluid ounces per acre (0.176 lb. ai/A) as a transplant tray drench no earlier than 72 hours prior to planting in the field.

SPRAY PREPARATION

Adjusting the pH of the application solution: The pH of the application solution should be between pH 4 and 6. (See "Acidification of Application Solution" section of the container label.)

SOIL APPLICATION DIRECTIONS

VERIMARK insect control may be applied as a transplant tray drench by growers or commercial transplant producers no earlier than 72 hours prior to planting in the field.

Refer to the SOIL APPLICATIONS section of the container label for additional use and application guidance.

Read and follow all label directions for the tray drench application method as described on the Federal Label.

See the RESISTANCE MANAGEMENT section of the VERIMARK insect control Federal label for additional guidance regarding insecticide resistance management.

See the TANK MIXTURES AND CROP SAFETY section of the VERIMARK insect control label for more information and for guidance regarding crop response for tested tank mixtures.

USE RESTRICTIONS

Do not apply more than 13.5 fluid ounces per acre (0.176 lb. ai/A) of VERIMARK insect control at planting.

Do not apply a total of more than 0.4 lb. ai/A of CYAZYPYR® active or cyantranilprole containing products per calendar year whether the application is made to the soil or foliarly.

Re-entry interval (REI) is 4 hours.

Pre-harvest interval (PHI) is 1 day.

R-4932 020525 021325



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000

FMC, the FMC logo, Cyazapyr and Verimark
are trademarks of FMC Corporation or an affiliate.
©2025 FMC Corporation