



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9606

FOR DISTRIBUTION AND USE ONLY IN DELAWARE, MAINE, MARYLAND, MICHIGAN, NEW JERSEY, NORTH CAROLINA, PENNSYLVANIA AND VIRGINIA

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation for CORAGEN® insect control is valid until December 31, 2025, or until withdrawn, cancelled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

REDUCED RATE APPLICATION FOR CONTROL OF CABBAGE LOOPER LARVAE IN POTATOES

Directions For Use

PEST CONTROLLED

CORAGEN insect control is recommended for the control of Cabbage Looper larvae in potatoes.

APPLICATION RATES

Apply CORAGEN insect control at 2.5 – 3.5 fluid ounces per acre as a foliar spray or via overhead chemigation. For best results with foliar sprays, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v). For best results via overhead chemigation applications, apply in 0.1- to 0.2-acre inches of water and add MSO at 12 - 16 fluid ounces per acre.

CABBAGE LOOPER APPLICATION TIMING

Apply when populations reach local established thresholds to prevent crop damage.



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104

CORAGEN insect control applied at the 2.5 – 3.5 fluid ounces per acre rate should be used on early-stage instars of Cabbage Looper (1st through 3rd instar). For larger larvae that may require higher rates (3.5 to 7.5 fluid ounces per acre), consult the EPA-registered label.

See the APPLICATION section of the CORAGEN insect control Federal label for additional guidance and restrictions regarding overhead chemigation.

See the RESISTANCE MANAGEMENT section of the CORAGEN insect control Federal label for additional guidance regarding insecticide resistance management.

USE RESTRICTIONS

Do not apply more than 15.4 fluid ounces of CORAGEN insect control or 0.2 lb a.i. of chlorantraniliprole-containing products per acre per calendar year.

Do not make more than 4 applications per acre per calendar year.

The minimum retreatment interval is 5 days.

The preharvest interval is 14 days.

The field re-entry interval is 4 hours.

Read and follow all applicable directions, restrictions, and precautions on the EPA-registered label.

IMPORTANT NOTICE

BEFORE BUYING OR USING FMC PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY, AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of Federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This Bulletin contains supplemental instructions for use of the referenced FMC product. Follow these instructions carefully. This Bulletin must be in the possession of the User at the time of application. **Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using the FMC product(s) covered by this Bulletin. FMC will not be responsible for losses or damages resulting from use of this product in any manner not specifically stated by FMC on the Section 3 label, or other labels, or this Bulletin. User assumes all risks associated with any such nonspecified use. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability of the Section 3 Federal product label**

R-4407 043020 04-29-20

FMC, the FMC logo and Coragen are trademarks of FMC Corporation or an affiliate.
© 2020 FMC Corporation



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104