

	<b>FIFRA 2(ee) Recommendation</b>
	EPA Reg. No. 279-9606
	<b>FOR DISTRIBUTION AND USE ONLY IN THE STATES OF IDAHO, OREGON AND WASHINGTON</b>
	<b>THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.</b>

This recommendation for CORAGEN® insect control is valid until December 31, 2025, or until withdrawn, cancelled or suspended.

<b>IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.</b>
<b>THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.</b>

## FOR CONTROL OF CERTAIN PESTS IN GRASS GROWN FOR SEED

Directions For Use
<p>CORAGEN insect control is recommended at 3.5 to 5.0 fluid ounces per acre (0.045 - 0.065 lb a.i. per acre) for the control of Armyworms, Cutworms, and Sod Webworms. For suppression of Billbug grubs and European Crane Fly larvae, apply CORAGEN insect control at 5.0 fluid ounces/acre (0.065 lb a.i. per acre).</p> <p><b>Method of Application</b></p> <p>For control of Armyworms and Cutworms, apply CORAGEN insect control as a foliar spray using properly calibrated ground equipment in a minimum of 10 gallons per acre, or via overhead chemigation in 0.10 to 0.20 acre inch of water.</p> <p>For foliar sprays, increase the spray volume to compensate for the amount of foliage present. For maximum spray penetration in to the root crown area, the use of a silicone surfactant may be useful. For best results with foliar spray applications, delay the next irrigation for at least 24 hours.</p> <p>For control of Sod Webworm larvae, and suppression of Billbug grubs and Crane Fly larvae, it is important to move the CORAGEN insect control into the grass root zone. This is best achieved by applying via overhead chemigation</p>



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104

in 0.25 to 0.50 acre inch of water, or by immediately following a foliar spray application with 0.25 to 0.50 acre inch of water.

#### **Application Timing**

For control of Armyworms, Cutworms, and Sod Webworms, apply at first sign of economic crop damage.

For suppression of European Crane Fly larvae, apply between September and early November.

For suppression of Billbug grubs, apply when overwintered adult Billbugs are first observed. This will usually occur in late April or early May.

**See the RESISTANCE MANAGEMENT section of the CORAGEN insect control Federal label for additional guidance regarding insecticide resistance management.**

#### **PRECAUTIONS**

Read and follow all instructions and precautions on the federal section 3 label before using CORAGEN insect control on grass grown for seed.

The preharvest interval is zero days.

The minimum re-entry interval is 4 hours.

Do not make more than 4 applications per crop.

The minimum interval between applications is 7 days.

Do not apply more than 15.4 fluid ounces per acre (0.2 lb a.i. per acre) per calendar year.

R-4493 121620 12-07-20

FMC, the FMC logo and Coragen are trademarks of FMC Corporation or an affiliate.

© 2020 FMC Corporation



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104