



## FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9606

**FOR DISTRIBUTION AND USE  
ONLY IN CALIFORNIA**

**THIS RECOMMENDATION, WHICH CONTAINS  
ADDITIONAL DIRECTIONS FOR USE, IS MADE  
AS PERMITTED UNDER FIFRA SECTION 2(ee)  
AND HAS NOT BEEN SUBMITTED TO OR  
APPROVED BY THE US EPA.**

This 2(ee) recommendation for CORAGEN insect control is valid until December 31, 2025, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

**THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT  
THE TIME OF PESTICIDE APPLICATION.**

### **Sugar Beets: Control of Beet Armyworm (*Spodoptera exigua*) and Western Yellow Striped Armyworm (*Spodoptera praefica*) by Soil Shank Injection of CORAGEN insect control**

#### **DIRECTIONS FOR USE**

Coragen® insect control is recommended for control of beet armyworm (*Spodoptera exigua*) and western yellow striped armyworm (*Spodoptera praefica*) by soil shank injection in sugar beets.

Apply Coragen® insect control at 3.5 – 7.5 fluid ounces per acre (0.045-0.098 lb. a.i./A) as a shank injection at planting.

For best results, apply Coragen® insect control in a manner that ensures the application is in the sugar beet root zone. Incorporate the application using sufficient irrigation (typically 0.5 to 1.0 inch of water) to reach the root zone. The shank injection should be placed in the seed row or just below the seed line, within 1-2 inches of the seed line.



FMC Corporation  
2929 Walnut Street  
Philadelphia, PA 19104  
215-299-6000

Apply higher rates within the listed rate range when pests are expected to occur later in the crop growth cycle, when heavy infestations of beet armyworms or western yellow striped armyworms occur, or when extreme environmental conditions, such as rainy weather and high temperatures, are expected.

- Do not apply Coragen® insect control within 1 day of harvest.
- Do not enter, or allow worker entry, into treated areas during the restricted-entry (REI) of 4 hours.
- Make no more than 4 applications of Coragen® insect control per acre per crop.
- Do not apply more than 15.4 fluid ounces of Coragen® insect control, or 0.2 lb. a.i. of chlorantraniliprole containing products, per acre per crop.
- Do not apply more than 61.6 fluid ounces of Coragen® insect control, or 0.8 lb a.i. of chlorantraniliprole containing products, per acre per calendar year.

#### **IMPORTANT NOTICE**

**BEFORE BUYING OR USING FMC PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY, AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.**

It is a violation of Federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This Bulletin contains supplemental instructions for use of the referenced FMC product. Follow these instructions carefully. This Bulletin must be in the possession of the User at the time of application. **Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using the FMC product(s) covered by this Bulletin. FMC will not be responsible for losses or damages resulting from use of this product in any manner not specifically stated by FMC on the Section 3 label, or other labels, or this Bulletin. User assumes all risks associated with any such nonspecified use. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability of the Section 3 Federal product label.**

FMC and CORAGEN are trademarks of FMC Corporation or an affiliate.

R-4293 110724 11-07-24