

**THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee)  
AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.**



**EPA Reg. No. 279-9612**

**Reduced rate applications for control  
of Western bean cutworm in corn (field)  
and corn (pop).**

*THIS 2(ee) RECOMMENDATION  
EXPIRES ON MARCH 31, 2024 AND  
MUST NOT BE USED AFTER THIS DATE*

## **2(ee) Recommendation**

**FOR DISTRIBUTION AND USE  
ONLY IN: KANSAS, OKLAHOMA,  
COLORADO, WYOMING, NEW  
MEXICO, AND NEBRASKA**

**IT IS A VIOLATION OF FEDERAL LAW  
TO USE THIS PRODUCT IN A MANNER  
INCONSISTENT WITH ITS LABELING**

### **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

PREVATHON® insect control is recommended for control of Western bean cutworm in corn (field) and corn (pop). Apply PREVATHON® insect control at 10 fluid ounces per acre (0.034 lb ai/acre) as a foliar spray.

Make applications from early tassel emergence through silking as indicated by moth trap numbers and/or field scouting for egg masses/larvae. Follow local university recommendations for treatment thresholds and best timing.

Western bean cutworm larvae must be controlled before entering the ear. Young larvae typically move up to the tassel after hatching to feed, then back down the plant before entering the ear. Larvae feeding on foliage treated with PREVATHON® insect control will cease feeding within a few hours and die within a few days.

Thorough spray coverage is essential for best performance. Use a minimum of 10 gallons by ground or 2 gallons by air. Higher spray volume of 15 gal. by ground and 3 gal. by air will improve control.

PREVATHON® insect control may be mixed with approved agricultural adjuvants in situations where spray coverage is difficult to achieve such as closed canopy or dense foliage. An oil based adjuvant may improve deposition in hot dry weather. Follow directions on adjuvant manufacturer's label.

Chemigation: Refer to the PREVATHON® insect control Section 3 label for specific directions and precautions for application via chemigation.



FMC Corporation  
2929 Walnut Street  
Philadelphia, PA 19104

## RESTRICTIONS AND PRECAUTIONS

Make no more than four (4) PREVATHON® insect control applications per acre per calendar year.

Avoid using the same mode of action (IRAC group number) on consecutive generations of insect pests.

Within the “Group 28 treatment window”, make no more than 2 applications of PREVATHON® or other Group 28 insecticides within a single generation of the target pest on a crop or within a 30 day period to the same insect species on a crop.

Minimum interval between treatments is 7 days.

Do not apply more than 60 fl. Oz. PREVATHON® insect control, or 0.2 lb a.i. of Chlorantraniliprole containing products per acre per calendar year.

The preharvest interval is 14 days for ears, and 1 day for forage, fodder, silage, and stover.

The field re-entry interval is 4 hours.

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED PREVATHON INSECT CONTROL LABEL (EPA REG. NO. 279-9612) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.** FMC will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by FMC. User assumes all risk associated with such non-directed use.

### IMPORTANT NOTICE

**BEFORE BUYING OR USING FMC PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY, AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.**

It is a violation of Federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This Bulletin contains supplemental instructions for use of the referenced FMC product. Follow these instructions carefully. This Bulletin must be in the possession of the User at the time of application. **Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using the FMC product(s) covered by this Bulletin. FMC will not be responsible for losses or damages resulting from use of this product in any manner not specifically stated by FMC on the Section 3 label, or other labels, or this Bulletin. User assumes all risks associated with any such nonspecified use. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability of the Section 3 Federal product label.**

FMC and PREVATHON® are trademarks or registered trademarks of FMC Corporation or an affiliate.