



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9612

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF ARKANSAS, DELAWARE, KANSAS, KENTUCKY, LOUISIANA, MARYLAND, MISSISSIPPI, MISSOURI, NEBRASKA, OKLAHOMA, PENNSYLVANIA, TENNESSEE, AND VIRGINIA

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation for PREVATHON® insect control is valid until December 31, 2025, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR CONTROL OF DECTES STEM BORER IN SOYBEAN

Directions For Use

PREVATHON insect control is recommended for control of Dectes stem borer (*Dectes texanus*) in soybeans.

Apply PREVATHON insect control at 20 fluid ounces per acre (0.067 lb ai/acre) as a foliar spray. As an alternative, apply PREVATHON insect control at 14 fluid ounces per acre (0.047 lb ai/acre) in a tank mix with STEWARD EC insecticide (EPA Reg. No. 279-9596) at 6 fluid ounces per acre (0.059 lb ai/acre) as a foliar spray. Use a minimum of 10 gallons per acre (GPA) of water for ground applications or 3 GPA for aerial applications. For best results, use 20 GPA by ground or 5 GPA by air. See product labels for chemigation instructions.

To minimize crop damage by the pest, apply at the onset of adult beetle flight. Ensure thorough spray coverage and make application to soybeans prior to egg laying. For best results, regular scouting using a sweep net is necessary



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000

to identify the emergence and infestation of adult beetles. If regular scouting is not used, apply at 1500 Growing Degree Days (GDD) in Nebraska and northern Kansas or consult with your local agricultural advisor for advice on application timing. Continued scouting should be used to track the duration of the emergence period. A second application may be necessary at 3 to 4 weeks after the initial application if adults continue to emerge over an extended period.

See the RESISTANCE MANAGEMENT section of the PREVATHON insect control Section 3 Federal label for additional guidance on insecticide resistance management.

Use Restrictions

Do not apply more than 60 fluid ounces of PREVATHON insect control or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.

Do not apply more than 45 fluid ounces of STEWARD EC insecticide or 0.44 lb a.i. of indoxacarb containing products per acre per calendar year.

Do not make more than 4 applications per acre per calendar year.

For PREVATHON insect control, the minimum interval between treatments is 3 days and preharvest interval is 1 day. If STEWARD EC insecticide is used, the minimum interval between treatments is 5 days and preharvest interval is 21 days.

For PREVATHON insect control the field re-entry interval is 4 hours. If STEWARD EC insecticide is used, the field re-entry interval is 12 hours.

R-4497 110320 12-07-20

FMC, the FMC logo and PREVATHON and STEWARD are Trademarks of FMC Corporation or an affiliate.

©2020 FMC Corporation



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000