SUPPLEMENTAL LABELING FOR USE ON RICE



WITH THE ACTIVE INGREDIENT RYNAXYPYR®

EPA Reg. No. 279-9612

This Supplemental Label expires on October 27, 2023 and must not be used or distributed after this date.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.

This Supplemental Labeling must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for PREVATHON® before applying. Carefully follow all precautionary statements and application use directions.

Crop	Application Method	Target Pest		fluid ounces product per	Last Application (Days to Harvest)	REI (Hours)
\	SOIL APPLICATION† Broadcast spray		0.08 – 0.1	24 - 30	N/A	4

USE RESTRICTIONS

Apply higher rates within the listed range for heavier infestations, larger/denser crops or extreme environmental conditions such as rainy weather and high temperatures.

† Only for application as a broadcast spray to soil: For water-seeded rice, apply PREVATHON to soil surface prior to seeding and flooding. For dry-seed rice, PREVATHON may be applied to the surface of the soil before, during or after planting, but application must be made before rice emergence.

Do not apply more than 5 days prior to flooding.

Once flood is established, hold the water for a minimum of 14 days before discharging the water.

Broadcast application may be made using aerial or ground application equipment.

Do not apply more than 30 fl oz PREVATHON® or 0.1 lb a.i. of chlorantraniliprole containing products per acre per calendar year.

Do not use PREVATHON® treated rice fields for the aquaculture of edible fish or crustacea (including crawfish) during the rice production cycle (planting through harvest).

R-4363A 011121 10-27-20

