



THIFENSULFURON METHYL	GROUP	2	HERBICIDE
TRIBENURON METHYL	GROUP	2	HERBICIDE

## Soluble Granule

For Use on Wheat (including durum), Barley, Oats, Triticale and Fallow

Active Ingredients	By Weight
Thifensulfuron-methyl Methyl 3-[[[4-methoxy-6-methyl-1,3,5-triazin-2-yl) amino]carbonyl]amino)sulfonyl]-2-thiophenecarboxylate	25%
Tribenuron-methyl Methyl 2-[[[N-(4-methoxy-6-methyl-1,3,5-triazin-2-yl)methylamino]carbonyl]amino)sulfonyl]benzoate	25%
Other Ingredients	50%
TOTAL	100.0%

Contains 0.25 lb Thifensulfuron Methyl per pound

Contains 0.25 lb Tribenuron Methyl per pound

EPA Reg. No. 279-9601

EPA Est. No. \_\_\_\_\_

**Nonrefillable Container**

**Refillable Container**

Net: \_\_\_\_\_ OR

Net: \_\_\_\_\_

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

### FIRST AID

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION!** Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

For medical emergencies involving this product, call toll free 1-800-331-3148.

Sold By



FMC Corporation  
2929 Walnut Street  
Philadelphia, PA 19104

## PERSONAL PROTECTIVE EQUIPMENT (PPE)

### Applicators and other handlers must wear:

Long-sleeved shirt and long pants.  
Chemical Resistant Gloves  $\geq$  14 mls.  
Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## USER SAFETY RECOMMENDATIONS

**USERS SHOULD:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

### Groundwater Advisory

This product has properties and characteristics associated with chemicals detected in groundwater. This product may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

### Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several weeks or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this product from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

### Windblown Soil Particles Advisory

This product has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying this product if prevailing local conditions may be expected to result in off-site movement.

### Non-target Organism Advisory

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift. For further guidance and instructions on how to minimize spray drift, refer to the Spray Drift Management section of this label.

## PRODUCT INFORMATION

AFFINITY® BroadSpec herbicide (with TotalSol® soluble granules) – referred to below as AFFINITY BroadSpec herbicide - is a soluble granule that is used for selective postemergence weed control in wheat (including durum), barley, oats, triticale and fallow. The best control is obtained when AFFINITY BroadSpec herbicide is applied to young, actively growing weeds. The use rate will depend on weed spectrum and size of weed at time of application. The degree and duration of control may depend on the following:

- weed spectrum and infestation intensity
- weed size at application
- environmental conditions at and following treatment

AFFINITY BroadSpec herbicide is noncorrosive, nonflammable, nonvolatile, and does not freeze. AFFINITY BroadSpec herbicide must be mixed, and completely dissolved in water and applied as a uniform broadcast spray.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AFFINITY BroadSpec herbicide must be used only in accordance with instructions on this label. To the extent consistent with applicable law, FMC will not be responsible for losses or damage resulting from the use of this product in any manner not specifically specified by FMC.

AFFINITY BroadSpec herbicide is for use on wheat (including durum), barley, oats, triticale and fallow in many states. Check with your state extension or Dept. of Agriculture before use, to be certain AFFINITY BroadSpec herbicide is registered in your state.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

Coveralls.

Chemical-resistant gloves made of any waterproof material.

Shoes plus socks.

### RESTRICTIONS

Injury to or loss of adjacent sensitive crops, desirable trees or vegetation may result from failure to observe the following:

- DO NOT apply to wheat, barley, oats, and triticale underseeded with another crop.
- DO NOT harvest wheat or barley sooner than 45 days after the last application of AFFINITY BroadSpec herbicide.
- DO NOT apply, drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- DO NOT use on lawns, walks, driveways, or tennis courts.
- DO NOT apply this product through any type of irrigation system.
- DO NOT apply AFFINITY BroadSpec herbicide by air in the state of New York.

### PRECAUTIONS

Injury to or loss of adjacent sensitive crops, desirable trees or vegetation may result from failure to observe the following:

- Take all necessary precautions to avoid all direct or indirect contact (including spray drift) with non-target plants or areas.

Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than wheat, barley, oats, and triticale.

Dry, dusty field conditions may result in reduced control in wheel track areas.

AFFINITY BroadSpec herbicide must not be applied to wheat, barley, oats, and triticale that is stressed by severe weather conditions, drought (including low levels of subsoil moisture), low fertility, water-saturated soil, disease, or insect damage, as crop injury may result. Risk of injury is greatest when crop is in the 2 to 5- leaf stage. Severe winter stress, drought, disease, or insect damage following application also may result in crop injury.

Wheat, barley, oats, and triticale may differ in their response to various herbicides. FMC advises that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of AFFINITY BroadSpec herbicide to a small area.

Under certain conditions, including heavy rainfall, prolonged cold weather (daily high temperature less than 50° F), or wide fluctuations in day/night temperatures prior to or soon after AFFINITY BroadSpec herbicide application, temporary discoloration and/or crop injury may occur. To reduce the potential of crop injury, tank mix AFFINITY BroadSpec herbicide with 2,4-D (ester formulations perform best—see “Tank Mixtures” section of this label) and apply after the crop is in the tillering stage of growth.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.).

## ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

AFFINITY BroadSpec herbicide is absorbed primarily through the foliage of plants, rapidly inhibiting the growth of susceptible weeds. One to 3 weeks after application to weeds, leaves of susceptible plants appear chlorotic, and the growing point subsequently dies.

AFFINITY BroadSpec herbicide provides the best control in vigorously growing crops that shade competitive weeds. Weed control in areas of thin crop stand or seeding skips may not be as satisfactory. However, a crop canopy that is too dense at application can intercept spray and reduce weed control.

The herbicidal action of AFFINITY BroadSpec herbicide may be affected in crops stressed from adverse environmental conditions (including extreme temperatures or moisture), abnormal soil conditions, cultural practices, or variations in crop variety. In warm, moist conditions, the expression of herbicide symptoms is accelerated; in cold, dry conditions, expression of herbicide symptoms is delayed. In addition, weeds hardened-off by drought stress are less susceptible to AFFINITY BroadSpec herbicide.

## WEED RESISTANCE MANAGEMENT

AFFINITY BroadSpec herbicide, which contains the active ingredients Thifensulfuron methyl and Tribenuron methyl, is a group 2 herbicide based on the mode of action classification system of the Weed Science Society of America.

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different sites of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices.

Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistance.

The continued effectiveness of this product depends on the successful implementation of a weed resistance management program.

To aid in the prevention of developing weeds resistant to this product, users should:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small (less than 4 inches).
- Apply full rates of AFFINITY BroadSpec herbicide for the most difficult to control weed in the field at the specified time (correct weed size) to minimize weed escapes.
- Scout fields after application to detect weed escapes or shifts in control of weed species.
- Control weed escapes before they reproduce by seed or proliferate vegetatively.
- Report any incidence of non-performance of this product against a particular weed to your FMC representative, local retailer, or county extension agent.
- Contact your FMC representative, crop advisor, or extension agent to find out if suspected resistant weeds to these MOAs have been found in your region. DO NOT assume that each listed weed is being controlled by multiple sites of action. Products with multiple active ingredients are intended to broaden the spectrum of weeds that are controlled. Some weeds may be controlled by only one of the active ingredient in this product.
- If resistance is suspected, treat weed escapes with an herbicide having a site of action other than Group 2 and/or use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production.
- Suspected herbicide-resistant weeds may be identified by these indicators:
  - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
  - A spreading patch of non-controlled plants of a particular weed species; and
  - Surviving plants mixed with controlled individuals of the same species.

Additionally, users should follow as many of the following herbicide resistance management practices as is practical:

- Use a broad spectrum soil-applied herbicide with other sites of action as a foundation in a weed control program.
- Utilize sequential applications of herbicides with alternative sites of action.
- Rotate the use of this product with non-Group 2 herbicides.
- Avoid making more than two applications of AFFINITY BroadSpec herbicide and any other Group 2 herbicides within a single growing season unless mixed with an herbicide with a different site of action with an overlapping spectrum for the difficult-to-control weeds.
- Incorporate non-chemical weed control practices, including mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.
- Use good agronomic principles that enhance crop development and crop competitiveness.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.
- Manage weeds in and around fields, during and after harvest to reduce weed seed production.

## INTEGRATED PEST MANAGEMENT

FMC advises the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Base the application of this product on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

### RATE CONVERSION CHART FOR AFFINITY BROADSPEC HERBICIDE

Ounces of AFFINITY BroadSpec herbicide/A	Pounds of AFFINITY BroadSpec herbicide /A	Active Ingredient	Pounds of Active Ingredient/A
0.4	0.025	Thifensulfuron methyl	0.0063
		Tribenuron methyl	0.0063
0.5	0.0313	Thifensulfuron methyl	0.0078
		Tribenuron methyl	0.0078
0.6	0.0375	Thifensulfuron methyl	0.0094
		Tribenuron methyl	0.0094
0.8	0.05	Thifensulfuron methyl	0.0125
		Tribenuron methyl	0.0125
1.0	0.0625	Thifensulfuron methyl	0.0156
		Tribenuron methyl	0.0156

## LABELLED USES

AFFINITY BroadSpec herbicide provides selective postemergence control of certain broadleaf weeds in wheat (including durum), barley, oat (spring and winter), triticale, post-harvest burndown, pre-plant burndown and fallow.

### Wheat (Including Durum), Barley, and Triticale

Application and Use Rate Information	Use Rates (oz of AFFINITY BroadSpec herbicide per acre)	Active Ingredient	Pounds of Active Ingredient per acre
Make applications after the crop is in the 2-leaf stage, but before the flag leaf is visible. Apply 0.4 to 1.0 oz AFFINITY BroadSpec herbicide per acre. Sequential treatments of AFFINITY BroadSpec herbicide may be made provided the total amount of AFFINITY BroadSpec herbicide applied to the crop per year does not exceed 1.0 oz/A.	0.4 to 1.0	Thifensulfuron methyl	0.0063 to 0.0156
		Tribenuron methyl	0.0063 to 0.0156

#### ***RESTRICTIONS in Wheat (including durum), Barley, and Triticale:***

- DO NOT apply to wheat, barley, or triticale crops underseeded with another crop.
- DO NOT harvest wheat or barley sooner than 45 days after the last application of AFFINITY BroadSpec herbicide.
- DO NOT apply more than 1.0 oz of AFFINITY BroadSpec herbicide per acre in a single application (maximum active ingredient per single application is 0.0156 lb/A thifensulfuron methyl and 0.0156 lb/A tribenuron methyl).
- DO NOT exceed two applications of AFFINITY BroadSpec herbicide per year in Wheat (including durum), Barley and Triticale when using reduced application rates.
- DO NOT apply more than 1.0 oz of AFFINITY BroadSpec herbicide per acre per year (maximum active ingredient load per year is 0.0156 lb/A thifensulfuron methyl and 0.0156 lb/A tribenuron methyl).
- When applying 0.4 to 0.6 oz per acre, AFFINITY BroadSpec herbicide must be used in a tank-mix combination with other registered herbicides.
- The Minimum Retreatment Interval is 14 days.
- PHI is 7 days for forage, 30 days for hay, and 45 days for wheat, barley and triticale.

#### ***PRECAUTIONS in Wheat (including durum), Barley, and Triticale:***

Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than wheat, barley, and triticale.

AFFINITY BroadSpec herbicide must not be applied to wheat, barley, and triticale that is stressed by severe weather conditions, drought (including low levels of subsoil moisture), low fertility, water-saturated soil, disease, or insect damage, as crop injury may result. Risk of injury is greatest when crop is in the 2 to 5- leaf stage. Severe winter stress, drought, disease, or insect damage following application also may result in crop injury.

Wheat, barley, and triticale may differ in their response to various herbicides. FMC suggests that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of AFFINITY BroadSpec herbicide to a small area.

#### ***TANK MIXTURES in Wheat (including durum), Barley, and Triticale:***

For expanded weed control, AFFINITY BroadSpec herbicide may be tank mixed with approved labeled rates of other herbicides labeled for use in wheat (including durum), barley and triticale. Refer to the other product's label for rotational crop intervals and other directions for use. When applying 0.4 to 0.6 oz/A in wheat, barley and triticale, AFFINITY BroadSpec herbicide must be used in a tank-mix combination with other registered herbicides.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. If the instructions on the tank mix partner label conflict with this AFFINITY BroadSpec herbicide, then DO NOT use in a tank mixture with AFFINITY BroadSpec herbicide.

## Winter Oat

Application and Use Rate Information	Use Rates (oz of AFFINITY BroadSpec herbicide per acre)	Active Ingredient	Pounds of Active Ingredient per acre
Make applications after the crop is in the 2-leaf stage, but before the flag leaf is visible.  Apply 0.4 oz/A AFFINITY BroadSpec herbicide for control of light populations of the weeds listed in Weeds Controlled table.  In oat, AFFINITY BroadSpec herbicide must be used in a tank-mix combination with other registered herbicides.	0.4	Thifensulfuron methyl	0.0063
		Tribenuron methyl	0.0063
<b><i>RESTRICTIONS in Winter Oat:</i></b> <ul style="list-style-type: none"><li>• DO NOT apply to oat crops underseeded with another crop.</li><li>• DO NOT apply more than 0.4 oz/A of AFFINITY BroadSpec herbicide in a single application (maximum active ingredient per single application is 0.0063 lb/A thifensulfuron methyl and 0.0063 lb/A tribenuron methyl).</li><li>• DO NOT apply more than 0.4 oz/A of AFFINITY BroadSpec herbicide per year (maximum active ingredient load per year is 0.0063 lb/A thifensulfuron methyl and 0.0063 lb/A tribenuron methyl) in Winter Oat.</li><li>• DO NOT make more than one application (or more than 0.0063 lb/A of active ingredient tribenuron-methyl) of AFFINITY BroadSpec herbicide per year on Winter Oat.</li><li>• The REI is 12 hours.</li></ul>			

### PRECAUTIONS in Winter Oat:

Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than oat.

AFFINITY BroadSpec herbicide must not be applied to oat that is stressed by severe weather conditions, drought (including low levels of subsoil moisture), low fertility, water-saturated soil, disease, or insect damage, as crop injury may result.

Risk of injury is greatest when crop is in the 2 to 5-leaf stage. Severe winter stress, drought, disease, or insect damage following application also may result in crop injury.

Oat may differ in their response to various herbicides. FMC suggests that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of AFFINITY BroadSpec herbicide to a small area.

### TANK MIXTURES in Winter Oat:

For expanded weed control, AFFINITY BroadSpec herbicide must be tank mixed with approved labeled rates of other herbicides labeled for use in winter oat. Refer to the other product's label for rotational crop intervals and other directions for use.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. If the instructions on the tank mix partner label conflict with this AFFINITY BroadSpec herbicide, then DO NOT use in a tank mixture with AFFINITY BroadSpec herbicide.



## Spring Oat

Application and Use Rate Information	Use Rates (oz of AFFINITY BroadSpec herbicide per acre)	Active Ingredient	Pounds of Active Ingredient per acre
<p>Make applications after the crop is in the 3-leaf stage but before jointing.</p> <p>Apply 0.4 oz/A AFFINITY BroadSpec herbicide for control of light populations of the weeds listed in Weeds Controlled table.</p> <p>In oat, AFFINITY BroadSpec herbicide must be used in a tank-mix combination with other registered herbicides.</p>	0.4	Thifensulfuron methyl	0.0063
		Tribenuron methyl	0.0063

### RESTRICTIONS in Spring Oat:

- DO NOT apply to oat crops underseeded with another crop.
- DO NOT apply more than 0.4 oz/A of AFFINITY BroadSpec herbicide in a single application (maximum active ingredient per single application is 0.0063 lb/A thifensulfuron methyl and 0.0063 lb/A tribenuron methyl).
- DO NOT apply more than 0.4 oz/A of AFFINITY BroadSpec herbicide per year (maximum active ingredient load per year is 0.0063 lb/A thifensulfuron methyl and 0.0063 lb/A tribenuron methyl) in Spring Oat.
- DO NOT make more than one application (or more than 0.0063 lb/A of active ingredient tribenuron-methyl) of AFFINITY BroadSpec herbicide per year on Spring Oat.
- The REI is 12 hours.
- DO NOT use on "Ogle", "Porter" or "Premier" varieties as crop injury can occur.

### PRECAUTIONS in Spring Oat:

Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than oat.

AFFINITY BroadSpec herbicide must not be applied to oat that is stressed by severe weather conditions, drought (including low levels of subsoil moisture), low fertility, water-saturated soil, disease, or insect damage, as crop injury may result. Risk of injury is greatest when crop is in the 2 to 5-leaf stage. Severe winter stress, drought, disease, or insect damage following application also may result in crop injury.

Oat may differ in their response to various herbicides. FMC suggests that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of AFFINITY BroadSpec herbicide to a small area.

### TANK MIXTURES in Spring Oat:

For expanded weed control, AFFINITY BroadSpec herbicide must be tank mixed with approved labeled rates of other herbicides labeled for use in spring oat. Refer to the other product's label for rotational crop intervals and other directions for use.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. If the instructions on the tank mix partner label conflict with this AFFINITY BroadSpec herbicide, then DO NOT use in a tank mixture with AFFINITY BroadSpec herbicide.



### ***Pre-Plant Burndown***

Application and Use Rate Information	Use Rates (oz of AFFINITY BroadSpec herbicide per acre)	Active Ingredient	Pounds of Active Ingredient per acre
<p>Apply AFFINITY BroadSpec herbicide as a burndown treatment to wheat (including durum) and barley to control emerged weeds prior to, or shortly after planting (prior to emergence). Make applications when the majority of weeds have emerged and are actively growing.</p> <p>AFFINITY BroadSpec herbicide can be used as a burndown treatment prior to planting other crops. See “CROP ROTATION” for the time interval required before planting.</p> <p>Apply 0.4 to 1.0 oz/A AFFINITY BroadSpec herbicide as a burndown treatment prior to, or shortly after planting (prior to emergence).</p> <p>When applying 0.4 to 0.6 oz/A, AFFINITY BroadSpec herbicide must be used in a tank-mix combination with other registered herbicides.</p> <p>Sequential treatments of AFFINITY BroadSpec herbicide may also be made provided the total amount of AFFINITY BroadSpec herbicide applied during one year does not exceed 1.0 oz/A.</p>	0.4 to 1.0	Thifensulfuron methyl	0.0063 to 0.0156
		Tribenuron methyl	0.0063 to 0.0156
<b><i>RESTRICTIONS for Pre-Plant Burndown (Wheat including durum, Barley, and other crops):</i></b> <ul style="list-style-type: none"><li>• DO NOT use less than 0.4 oz/A AFFINITY BroadSpec herbicide, unless otherwise specified by FMC.</li><li>• DO NOT apply more than 1.0 oz/A of AFFINITY BroadSpec herbicide in a single application (maximum active ingredient per single application is 0.0156 lb/A thifensulfuron methyl and 0.0156 lb/A tribenuron methyl).</li><li>• DO NOT apply more than 1.0 oz/A of AFFINITY BroadSpec herbicide per year (maximum active ingredient load per year is 0.0156 lb/A thifensulfuron methyl and 0.0156 lb/A tribenuron methyl).</li><li>• DO NOT exceed two applications of AFFINITY BroadSpec herbicide per year for Pre-Plant Burndown (Wheat including durum, Barley, and other crops) when using reduced application rates.</li><li>• The Minimum Retreatment Interval is 14 days.</li></ul>			

### ***TANK MIXTURES for Pre-Plant Burndown:***

AFFINITY BroadSpec herbicide may be used as a pre-plant burndown treatment alone or tank mixed with approved labeled rates of other herbicides labeled for use as a pre-plant burndown product. Refer to the other product's label for rotational crop intervals and other directions for use. When applying 0.4 to 0.6 oz/A as a pre-plant burndown treatment, AFFINITY BroadSpec herbicide must be used in a tank-mix combination with other registered herbicides.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. If the instructions on the tank mix partner label conflict with this AFFINITY BroadSpec herbicide, then DO NOT use in a tank mixture with AFFINITY BroadSpec herbicide.

## **Post-Harvest Burndown**

<b>Application and Use Rate Information</b>	<b>Use Rates (oz of AFFINITY BroadSpec herbicide per acre)</b>	<b>Active Ingredient</b>	<b>Pounds of Active Ingredient per acre</b>
<p>AFFINITY BroadSpec herbicide may be used as a burndown treatment to crop stubble when the majority of weeds have emerged and are actively growing. (See the "CROP ROTATION" section of this label for additional information).</p> <p>Apply AFFINITY BroadSpec herbicide at 0.4 to 1.0 oz/A to crop stubble after harvest. Use the 1.0 oz/A rate when weed infestation is heavy and predominantly consists of those weeds listed under the "WEEDS PARTIALLY CONTROLLED" section of this label or when application timing and environmental conditions are marginal. Apply AFFINITY BroadSpec herbicide in combination with other suitable registered burndown herbicides.</p> <p>Sequential treatments of AFFINITY BroadSpec herbicide may also be made provided the total amount of AFFINITY BroadSpec herbicide applied in one year does not exceed 1.0 oz/A.</p>	0.4 to 1.0	Thifensulfuron methyl	0.0063 to 0.0156
		Tribenuron methyl	0.0063 to 0.0156

### **RESTRICTIONS for Post-Harvest:**

- DO NOT use less than 0.4 oz/A AFFINITY BroadSpec herbicide, unless otherwise specified by FMC.
- DO NOT apply more than 1.0 oz/A of AFFINITY BroadSpec herbicide in a single application (maximum active ingredient per single application is 0.0156 lb/A thifensulfuron methyl and 0.0156 lb/A tribenuron methyl).
- DO NOT apply more than 1.0 oz/A AFFINITY BroadSpec herbicide per year (maximum active ingredient load per year is 0.0156 lb/A thifensulfuron methyl and 0.0156 lb/A tribenuron methyl).
- DO NOT exceed two applications of AFFINITY BroadSpec herbicide per year for Post-Harvest Burndown when using reduced application rates.
- The Minimum Retreatment Interval is 14 days.

### **TANK MIXTURES for Post-Harvest Burndown:**

AFFINITY BroadSpec herbicide may be used as a post-harvest treatment to crop stubble and must be tank mixed with approved labeled rates of other herbicides labeled for use as a post-harvest burndown treatment. Refer to the other product's label for rotational crop intervals and other directions for use.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. If the instructions on the tank mix partner label conflict with this AFFINITY BroadSpec herbicide, then DO NOT use in a tank mixture with AFFINITY BroadSpec herbicide.

## ***Fallow***

Application and Use Rate Information	Use Rates (oz of AFFINITY BroadSpec herbicide per acre)	Active Ingredient	Pounds of Active Ingredient per acre
<p>AFFINITY BroadSpec herbicide may be used as a fallow treatment, in the spring, summer or fall when the majority of weeds have emerged and are actively growing.</p> <p>Apply 0.4 to 1.0 oz/A AFFINITY BroadSpec herbicide to fallow. The total amount of AFFINITY BroadSpec herbicide cannot exceed 1.0 oz/A per year.</p> <p>When AFFINITY BroadSpec herbicide is applied at a rate of 0.4 to 0.6 oz/A, AFFINITY BroadSpec herbicide must be used in a tank-mix combination with other registered fallow herbicides. In fallow, AFFINITY BroadSpec herbicide may be applied in combination with other suitable registered fallow herbicides including glyphosate plus 2,4-D (ester formulations work best) or glyphosate plus dicamba.</p>	0.4 to 1.0	Thifensulfuron methyl	0.0063 to 0.0156
		Tribenuron methyl	0.0063 to 0.0156
<b><i>RESTRICTIONS in Fallow:</i></b>			
<ul style="list-style-type: none"><li>• DO NOT use less than 0.4 oz/A AFFINITY BroadSpec herbicide, unless otherwise specified by FMC.</li><li>• DO NOT apply more than 1.0 oz/A AFFINITY BroadSpec herbicide in a single application (maximum active ingredient per single application is 0.0156 lb/A thifensulfuron methyl and 0.0156 lb/A tribenuron methyl).</li><li>• DO NOT apply more than 1.0 oz /A AFFINITY BroadSpec herbicide per year (maximum active ingredient load per year is 0.0156 lb/A thifensulfuron methyl and 0.0156 lb/A tribenuron methyl).</li><li>• DO NOT exceed two applications of AFFINITY BroadSpec herbicide per year in Fallow when using reduced application rates.</li><li>• The Minimum Retreatment Interval is 14 days.</li></ul>			

### ***TANK MIXTURES in Fallow:***

AFFINITY BroadSpec herbicide may be used as a fallow treatment and must be tank mixed with approved labeled rates of other herbicides labeled for use in fallow. Refer to the other product's label for rotational crop intervals and other directions for use.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. If the instructions on the tank mix partner label conflict with this AFFINITY BroadSpec herbicide, then DO NOT use in a tank mixture with AFFINITY BroadSpec herbicide.

## TANK MIXTURES

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. If the instructions on the tank mix partner label conflict with this AFFINITY BroadSpec herbicide, then DO NOT use in a tank mixture with AFFINITY BroadSpec herbicide.

AFFINITY BroadSpec herbicide may be tank mixed with one or more suitable registered herbicides to control weeds listed as suppressed, weeds resistant to AFFINITY BroadSpec herbicide or weeds not listed under **Weeds Controlled**. Read and follow all manufacturer's label instructions for the companion herbicide. If those instructions conflict with this label, DO NOT tank mix the herbicide with AFFINITY BroadSpec herbicide.

AFFINITY BroadSpec herbicide can also be mixed with registered fungicides, insecticides, or liquid fertilizer for use on wheat, barley, or fallow.

### ***With Other Grass Control Products***

AFFINITY BroadSpec herbicide can be tank mixed with grass control products. Antagonism generally does not occur. Under certain environmental conditions, however, antagonism to Group 1 graminicides can occur. This may be reduced by using the higher rate of the graminicide. FMC advises that you first consult your state experiment station, university, or extension agent, Agricultural dealer, or FMC representative as to the potential for antagonism before using the mixture. If no information is available, limit the initial use of AFFINITY BroadSpec herbicide and the grass product to a small area.

### ***With Insecticides or Fungicides***

AFFINITY BroadSpec herbicide may be tank mixed or used sequentially with insecticides (or fungicides) registered for use on cereal grains. However, under certain conditions (drought stress, or if the crop is in the 2-4 leaf stage), tank mixes or sequential applications of AFFINITY BroadSpec herbicide with organophosphate insecticides (including parathion) may produce temporary crop yellowing or, in severe cases, crop injury. Test these mixtures in a small area before treating large areas. However, review all insecticide and fungicide labels for restrictions.

**DO NOT use AFFINITY BroadSpec herbicide plus products containing malathion, as crop injury will result.**

### ***With Liquid Nitrogen Solution Fertilizer***

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing AFFINITY BroadSpec herbicide in fertilizer solution. DO NOT add AFFINITY BroadSpec herbicide directly to liquid nitrogen fertilizer; the granules will not dissolve. AFFINITY BroadSpec herbicide must be thoroughly mixed with clean water before it is added to liquid nitrogen fertilizer. If granules remain when the mixture is poured out, add more clean water and mix until all granules have disappeared. Ensure that the agitator is running when the AFFINITY BroadSpec herbicide premix is added. Use of this mixture may result in temporary crop yellowing and stunting.

If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume), the addition of surfactant is necessary. Add surfactant at 1/4 qt - 1 qt per 100 gal of spray solution (0.06 -0.25% v/v) based on local guidance.

When using high rates of liquid nitrogen fertilizer solution in the spray solution, adding surfactant increases the risk of crop injury. Consult your agricultural dealer, consultant, fieldman, or FMC representative for specific instructions before adding an adjuvant to these tank mixtures.

If 2,4-D or MCPA is included with AFFINITY BroadSpec herbicide and fertilizer mixture, ester formulations tend to be more compatible (see manufacturer's label). Additional surfactant is not needed when using AFFINITY BroadSpec herbicide in tank mix with 2,4-D ester or MCPA ester and liquid nitrogen fertilizer solutions.

DO NOT use low rates of liquid nitrogen fertilizer solution as a substitute for a surfactant.

DO NOT use with liquid fertilizer solutions with a pH less than 3.0.

## CROP ROTATION

Labeled crops may be planted at specified time intervals following application of labeled rates of AFFINITY BroadSpec herbicide. Use the time intervals listed below to determine the required time interval before planting.

Refer to individual product labels to determine rotational crop restrictions when tank mixtures are used.

**Time Interval Before Planting\* (days after treatment with AFFINITY BroadSpec herbicide)**

<b>Crop</b>	<b>Days</b>
Barley, Rice, Soybeans with Bolt® technology, Triticale, and Wheat (including durum)	0
Soybeans	7†*
Cotton, Field Corn, and Grain Sorghum	14*
Sugarbeets, Winter Rape, and Canola	60
Any other crop	45

† When AFFINITY BroadSpec is applied at 0.5 oz/a or less the time interval for soybeans is 1 day.

\* Where AFFINITY BroadSpec is used on light textured soils, including sands and loamy sands, extend time to planting by 7 additional days. Where AFFINITY BroadSpec is used on high pH soils (>7.9), extend time to planting by 7 additional days.

## WEEDS CONTROLLED

AFFINITY BroadSpec herbicide effectively controls the following weeds when used according to label directions:

Annual knawel	London rocket
Annual sowthistle	Marshelder
Black mustard	Mayweed chamomile
Blue/Purple mustard	Miners lettuce
Broadleaf dock	Narrowleaf lambsquarters
Bur buttercup	Nightflowering catchfly
Bushy wallflower/Treacle mustard	Pennsylvania smartweed
Canada thistle *	Pineappleweed
Clasping pepperweed	Prickly lettuce*
Coast fiddleneck	Prostrate knotweed
Common buckwheat	Prostrate pigweed
Common chickweed	Redmaids
Common cocklebur *	Redroot pigweed
Common groundsel	Russian thistle*
Common lambsquarters	Scentless chamomile/mayweed
Common ragweed *	Shepherd's-purse
Common sunflower	Slimleaf lambsquarters
Corn chamomile	Smallflower buttercup
Corn gromwell *	Smallseed falseflax
Corn spurry	Stinking chickweed
Cowcockle	Stinking mayweed/dogfennel
Cress (mouse-ear)	Sunflower
Curly dock	Swinecress
False chamomile	Tansymustard
Field chickweed	Tarweed fiddleneck
Field pennycress	Tumble/Jim Hill mustard
Filaree (redstem, Texas)	Volunteer canola
Flixweed	Volunteer lentils
Green smartweed	Volunteer peas
Henbit	White cockle
Kochia *	Wild buckwheat*
Ladysthumb	Wild chamomile
Lanceleaf sage *	Wild mustard

## WEEDS PARTIALLY CONTROLLED\*\*

AFFINITY BroadSpec herbicide partially controls the following weeds when used according to label directions:

Catchweed bedstraw	Marestail
Mallow (common, little)	Nightshade (cutleaf, hairy)

\* See SPECIFIC WEED PROBLEMS for more information.

\*\* Partial control: A visual reduction of weed population as well as a significant loss of vigor. For better results, use the highest specified rate of AFFINITY BroadSpec herbicide per acre and include a tank mix partner including 2,4-D, MCPA, bromoxynil containing herbicides or dicamba containing herbicides.

## SPECIFIC WEED PROBLEMS

**Canada thistle:** For control in wheat and barley, use 0.8 oz/A AFFINITY BroadSpec herbicide plus surfactant when all thistles are 4" to 8" with 2" to 6" of new growth. Make the application in the spring. Control will be improved by using AFFINITY BroadSpec herbicide in combination with 2,4-D, dicamba, WideMatch® herbicide, or Colt® AS herbicide.

**Common cocklebur, Common ragweed, Lanceleaf sage:** In wheat and barley, apply AFFINITY BroadSpec herbicide at 0.4 to 0.8 oz/A in combination with 2,4-D at rates from 1/4 to 3/8 lb active ingredient (ester formulations work best) when weeds are small and actively growing. When using 1/4 lb active ingredient of 2,4-D, be sure to add surfactant at the rate of 1/4 to 1/2 quart per 100 gallons of spray solution (0.06 to 0.125% v/v—use the higher rate under stress conditions).

**Corn gromwell, Wild buckwheat:** For control in wheat and barley, use 0.8 oz/A AFFINITY BroadSpec herbicide plus surfactant.

**Kochia, Russian thistle, Prickly lettuce:** Naturally occurring resistant biotypes of these weeds are known to occur. For best results, use AFFINITY BroadSpec herbicide in a tank mix with Colt + Sword® herbicide, Colt + Salvo® herbicide, dicamba and 2,4-D; or Bromoxynil (including Buctril® herbicide) and 2,4-D (3/4 - 1 pt Buctril herbicide + 1/4 - 3/8 lb active ingredient 2, 4-D ester). For improved broadleaf weed control including kochia, AFFINITY BroadSpec herbicide can also be tank mixed with WideMatch® herbicide, Colt® AS herbicide or Starane® NXT herbicide. AFFINITY BroadSpec herbicide must be applied in the spring when weeds are 2" to 4" tall or 2" to 4" across and are actively growing. Refer to the Tank Mixtures section of this label for additional details.

**Yellow and Green Foxtail:** AFFINITY BroadSpec herbicide at 0.6 - 1.0 oz/A may be tank mixed with GoldSky® herbicide, "Everest®" branded products, or Rimfire® Max herbicide for yellow and green foxtail suppression in wheat.

## SPRAY ADJUVANTS

Always include a spray adjuvant with applications of AFFINITY BroadSpec herbicide. In addition to a spray adjuvant, an ammonium nitrogen fertilizer may be used.

Consult your Ag dealer or applicator, local FMC fact sheets, technical bulletins, and service policies prior to using an adjuvant system. If another herbicide is tank mixed with AFFINITY BroadSpec herbicide, select adjuvants authorized for use with both products. Products must contain only EPA-exempt ingredients (40CFR 1001).

### **Nonionic Surfactant (NIS)**

- Apply 0.06 to 0.50% volume/volume (1/2 pt to 4 pt per 100 gal of spray solution).
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12. See the Tank Mixtures section of this label for additional information.

### **Petroleum Crop Oil Concentrate (COC) or Modified Seed Oil (MSO)**

- Apply at 1% volume/volume (1 gal per 100 gal spray solution) or 2% volume/volume under arid conditions.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

### **Special Adjuvant Types**

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.
- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality and have been evaluated and approved by FMC product management.

### **Ammonium Nitrogen Fertilizer**

- Use 2 qt/acre of a high-quality urea ammonium nitrate (UAN), including 28%N or 32%N, or 2 lb/acre of a spray-grade ammonium sulfate (AMS). Use 4 qt/acre UAN or 4 lb/acre AMS under arid conditions.



## APPLICATION TIMING

Since AFFINITY BroadSpec herbicide has very little or no soil activity, it controls only those weeds that have germinated; therefore, apply AFFINITY BroadSpec herbicide when all or most of the weeds have germinated. Annual broadleaf weeds must be past the cotyledon stage, actively growing, and less than 4" tall or wide. Rainfall immediately after treatment can wash AFFINITY BroadSpec herbicide off of weed foliage, resulting in reduced weed control. Several hours of dry weather are needed to allow AFFINITY BroadSpec herbicide to be sufficiently absorbed by weed foliage.

## GROUND APPLICATION

For optimum spray distribution and thorough coverage, use flat-fan or low-volume flood nozzles.

For flat-fan nozzles, use a spray volume of at least 5 gal per acre (GPA).

For flood nozzles on 30" spacings, use at least 10 GPA, flood nozzles no larger than TK10 (or the equivalent), and a pressure of at least 30 psi. For 40" nozzle spacings, use at least 13 GPA; for 60" spacings use at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings.

Raindrop "RA" nozzles are not advised for AFFINITY BroadSpec herbicide applications, as weed control performance may be reduced.

Use screens that are 50-mesh or larger.

## AERIAL APPLICATION

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage at 2 to 5 GPA.

Use at least 3 GPA in Idaho, Oregon, or Utah.

This product is limited to ground application only in the State of New York.

See the **Spray Drift** section of this label.

## CHEMIGATION

DO NOT apply this product through any irrigation system.

## PRODUCT MEASUREMENT

AFFINITY BroadSpec herbicide is measured using the AFFINITY BroadSpec herbicide volumetric measuring cylinder. The degree of accuracy of this cylinder varies by  $\pm 7.5\%$ . For more precise measurement, use scales calibrated in ounces.

## PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Ensure that all operation employees accurately measure pesticides.
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- DO NOT discharge excess material on the soil at a single spot in the field, grove, or mixing/loading station.
- Dilute and agitate excess solution and apply at labeled rates or uses.
- Avoid storage of pesticides near well sites.
- When triple-rinsing the pesticide container, be sure to add the rinsate to the spray mix.

## MIXING INSTRUCTIONS

DO NOT use with spray additives that alter the pH of the spray solution below pH 5.0 or above pH 9.0 as rapid product degradation can occur. Spray solutions of pH 6.0-8.0 allow for optimum stability of AFFINITY BroadSpec herbicide.

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of AFFINITY BroadSpec herbicide.
3. Continue agitation until the AFFINITY BroadSpec herbicide is fully dissolved, at least 5 minutes.
4. Once the AFFINITY BroadSpec herbicide is fully dissolved, maintain agitation and continue filling tank with water.
5. As the tank is filling, add tank mix partners and then add the required volume of spray adjuvant. Always add spray adjuvant last. Antifoaming agents may be used. DO NOT use with spray additives that alter the pH of the spray solution below pH 6.0 as rapid product degradation can occur. Spray solutions of pH 7.0 and higher allow for optimum stability of AFFINITY BroadSpec herbicide.
6. Dispersed tank mix partners can settle if the tank mixture is not continually agitated. If settling occurs, thoroughly reagitate before using.
7. Apply AFFINITY BroadSpec herbicide spray mixture within 24 hours of mixing to avoid product degradation.
8. If AFFINITY BroadSpec herbicide and a tank mix partner are to be applied in multiple loads, fully dissolve the AFFINITY BroadSpec herbicide in clean water prior to adding to the tank.

## **GRAZING**

Allow at least 7 days between application and grazing of treated forage. In addition, allow at least 7 days between application and feeding of forage from treated areas to livestock. Allow at least 30 days between application and feeding of hay from treated areas to livestock. Harvested straw may be used for bedding and/or feed. Allow at least 45 days between application and harvesting of grain.

## **SPRAY EQUIPMENT**

For specific application equipment, refer to the manufacturer's instructions for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy, etc.

Be sure to calibrate air or ground equipment properly before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when crop canopy is dense. Avoid swath overlapping, and shut off spray booms while starting, turning, slowing, or stopping, to avoid injury to the crop. For additional information on spray drift refer to Spray Drift section of label.

Continuous agitation is not required for AFFINITY BroadSpec herbicide but may be required to keep tank-mix partners in solution or suspension. Refer to tank-mix partner labels for additional information.

## **SPRAYER CLEANUP**

The spray equipment must be cleaned before AFFINITY BroadSpec herbicide is sprayed. Follow the cleanup procedures specified on the labels of the previously applied products. If no directions are provided, follow the six steps outlined in "After Spraying AFFINITY BroadSpec herbicide" in this label.

## **AT THE END OF THE DAY**

It is advised that during periods when multiple loads of AFFINITY BroadSpec herbicide are applied, at the end of each day of spraying, the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits which can accumulate in the application equipment.

## **AFTER SPRAYING AFFINITY BROADSPEC HERBICIDE AND BEFORE SPRAYING CROPS OTHER THAN WHEAT AND BARLEY**

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of AFFINITY BroadSpec herbicide as follows:

1. Empty the tank and drain the sump completely.
2. Spray the tank walls with clean water using a minimum volume of 10% of the tank volume. Circulate the water through the lines, including all by-pass lines, for at least two minutes. Flush the boom well and empty the sprayer. Completely drain the sump.
3. Repeat step 2.
4. Remove the nozzles and screens and clean separately in a bucket containing water.

The rinsate solution may be applied to the crop(s) specified on this label. DO NOT exceed the maximum-labeled use rate. If cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

### **Notes:**

1. Always start with a clean spray tank.
2. Steam-cleaning aerial spray tanks is advised to facilitate the removal of any caked deposits.
3. When AFFINITY BroadSpec herbicide is tank mixed with other pesticides, all cleanout procedures for each product must be examined and the most rigorous procedure must be followed.
4. In addition to this cleanout procedure, all pre-cleanout guidelines on subsequently applied products must be followed as per the individual labels.

## MANDATORY SPRAY DRIFT

### Ground Boom Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy unless making a turf, pasture, or rangeland application, in which case applicators may apply with a nozzle height no more than 4 feet above the ground.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- DO NOT apply when wind speeds exceed 10 miles per hour at the application site.
- DO NOT apply during temperature inversions.

### Aerial Applications:

- DO NOT release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use one-half swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- DO NOT apply when wind speeds exceed 10 miles per hour at the application site.
- DO NOT apply during temperature inversions.

## SPRAY DRIFT MANAGEMENT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

## IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### Controlling Droplet Size – Aircraft

- Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles.
- Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

## **BOOM HEIGHT – GROUND BOOM**

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

## **RELEASE HEIGHT - AIRCRAFT**

Higher release heights increase the potential for spray drift. When applying aerially to crops, DO NOT release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

## **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

## **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

## **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

## **WIND**

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

## **Handheld Technology Applications:**

- Take precautions to minimize spray drift.

## **DRIFT CONTROL ADDITIVES**

Using product compatible drift control additives can reduce drift potential. When a drift control additive is used, read and carefully observe cautionary statements and all other information on the additive's label. If using an additive that increases viscosity, ensure that the nozzles and other application equipment will function properly with a viscous spray solution.

Preferred drift control additives have been certified by the Chemical Producers and Distributors Association (CPDA).

## FOR USE IN SEED PRODUCTION OF SOYBEANS WITH BOLT TECHNOLOGY

Application and Use Rate Information	Use Rates (oz of AFFINITY BroadSpec herbicide per acre)	Active Ingredient	Pounds of Active Ingredient per acre
<p>For control of broadleaf weeds and for the detection and control of susceptible soybean “segregates”, apply AFFINITY BroadSpec herbicide at 0.5 - 1.0 oz/A to soybeans with BOLT technology.</p> <p>Applications of AFFINITY BroadSpec herbicide on soybean may be made from emergence through the bloom growth stage. Best results are obtained when applications are made to actively growing plants.</p> <p>AFFINITY BroadSpec herbicide is rainfast 1 hour after application.</p>	0.5 to 1.0	Thifensulfuron methyl	0.0078 to 0.0156
		Tribenuron methyl	0.0078 to 0.0156

### RESTRICTIONS in seed production of soybeans with BOLT Technology:

- DO NOT apply more than 1.0 oz of AFFINITY BroadSpec herbicide per acre in a single application (maximum active ingredient per single application is 0.0156 lb/A thifensulfuron methyl and 0.0156 lb/A tribenuron methyl).
- DO NOT exceed two applications of AFFINITY BroadSpec herbicide per year in seed production of soybeans with BOLT Technology when using reduced application rates.
- DO NOT apply more than 1.0 oz of AFFINITY BroadSpec herbicide per acre per year (maximum active ingredient load per year is 0.0156 lb/A thifensulfuron methyl and 0.0156 lb/A tribenuron methyl).
- When applying 0.4 to 0.6 oz per acre, AFFINITY BroadSpec herbicide must be used in a tank-mix combination with other registered herbicides.
- The Minimum Retreatment Interval is 14 days.
- Soybeans containing the BOLT trait may be replanted anytime after the application of AFFINITY BroadSpec herbicide. Do not graze, harvest or feed forage or grain from treated soybeans within 7 days of application.

### PRECAUTIONS in seed production of soybeans with BOLT Technology:

Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than soybeans with BOLT technology.

AFFINITY BroadSpec herbicide is a water-soluble herbicide applied as a foliar spray to selectively control certain broadleaf weeds in soybeans with BOLT technology. AFFINITY BroadSpec herbicide may also be applied to remove susceptible “segregates”, i.e., undesirable soybean plants that do not contain the BOLT trait, during seed propagation. Soybean plants not possessing the BOLT trait will be severely injured or killed if treated with AFFINITY BroadSpec herbicide.

### **TANK MIXTURES in seed production of soybeans with BOLT Technology:**

When tank mixing AFFINITY BroadSpec herbicide + glyphosate herbicides, it is advised to add 4.25 -17 lb ammonium sulfate per 100 gallons of spray mixture. The addition of surfactant at 0.25% v/v (1 qt per 100 gallons of spray) to some AFFINITY BroadSpec herbicide + glyphosate tank mixes may result in improved weed control. See the glyphosate manufacturer's label for specific ammonium sulfate and surfactant directions.

Weed control and crop safety resulting from the use of tank mixtures not specifically noted on this label are the responsibility of the user.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. If the instructions on the tank mix partner label conflict with this AFFINITY BroadSpec herbicide, then DO NOT use in a tank mixture with AFFINITY BroadSpec herbicide.

### **WEEDS CONTROLLED**

In addition to detecting and controlling susceptible soybean "segregates", AFFINITY BroadSpec herbicide also effectively controls the following weeds when used according to label directions:

Annual knawel	Curly dock	Redmaids
Annual sowthistle	False chamomile	Redroot pigweed
Black mustard	Field chickweed	Russian thistle*
Blue/Purple mustard	Field pennycress	Scentless chamomile/mayweed
Broadleaf dock	Filaree (redstem, Texas)	Shepherd's-purse
Bur buttercup	Flixweed	Slimleaf lambsquarters
Bushy wallflower/Treacle mustard	Green smartweed	Smallflower buttercup
Canada thistle *	Henbit	Smallseed falseflax
Clasping pepperweed	Kochia *	Stinking chickweed
Coast fiddleneck	Ladysthumb	Stinking mayweed/dogfennel
Common buckwheat	Lanceleaf sage *	Sunflower
Common chickweed	London rocket	Swinecress
Common cocklebur *	Marshelder	Tansymustard
Common groundsel	Mayweed chamomile	Tarweed fiddleneck
Common lambsquarters	Miners lettuce	Tumble/Jim Hill mustard
Common ragweed *	Narrowleaf lambsquarters	Volunteer canola
Common sunflower	Nightflowering catchfly	Volunteer lentils
Corn chamomile	Pennsylvania smartweed	Volunteer peas
Corn gromwell *	Pineappleweed	White cockle
Corn spurry	Prickly lettuce*	Wild buckwheat*
Cowcockle	Prostrate knotweed	Wild chamomile
Cress (mouse-ear)	Prostrate pigweed	Wild mustard

\*These species may be difficult to control with, or, resistant to AFFINITY BroadSpec herbicide.

## IDENTIFICATION INFORMATION FOR PRODUCTS REFERENCED IN THIS LABEL

### REGISTERED PRODUCTS REFERENCED IN THIS LABEL FOR TANK MIXTURES OR MENTIONED FOR OTHER REASONS

Product Name	Active Ingredient(s)	EPA Registration Number
Colt® AS herbicide	Clopyralid + Fluroxypyr	34704-1019
Colt + Salvo® herbicide	Fluroxypyr + 2,4-D	34704-1010
Colt + Sword® herbicide	Fluroxypyr + MCPA	34704-1011
Buctril® herbicide	bromoxynil	264-437
Rimfire® Max herbicide	Mesosulfuron-methyl + Propoxycarbazone-sodium	264-1099
Everest® 2.0 herbicide	Flucarbazone-sodium	66330-391
Everest® 3.0 herbicide	Flucarbazone-sodium	66330-429
Everest® 3.0 AG	Flucarbazone-sodium	66330-433
WideMatch® herbicide	Clopyralid + Fluroxypyr	62719-512
Starane® NXT herbicide	Bromoxynil + Fluroxypyr	62719-557
GoldSky® herbicide	Fluroxypyr, 1-methylheptyl ester + Florasulam + Pyroxsulam	62719-582



## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store product in original container only. Store in a cool, dry place.

**Pesticide Disposal:** Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

**Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds):** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**Nonrefillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds):** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down):** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums With Liners:** Nonrefillable container. Do not reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

**Refillable Fiber Drums With Liners:** Refillable container (fiber drum only). *Refilling Fiber Drum:* Refill this fiber drum with AFFINITY BroadSpec herbicide containing thifensulfuron methyl and tribenuron methyl only. Do not reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. *Disposing of Fiber Drum and/or Liner:* Do not reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. Offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

**All Other Refillable Containers:** Refillable container. *Refilling Container:* Refill this container with AFFINITY BroadSpec herbicide containing thifensulfuron methyl and tribenuron methyl only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use the container, contact FMC at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, do not reuse or transport container, contact FMC at the number below for instructions. *Disposing of Container:* Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**Outer Foil Pouches of Water Soluble Packets (WSP):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or, dispose of the empty outer foil pouch in the trash as long as WSP is unbroken. If the outer pouch contacts the formulated product in any way, the pouch must be triple rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer pouch as described previously.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact CHEMTREC (Transportation and Spills) at 1-800-424-9300, day or night.

**NOTICE TO BUYER:** Purchase of this material does not confer any rights under patents of countries outside of the United States.

FMC, Affinity and TotalSol are trademarks of FMC Corporation or an affiliate.

Colt, Colt + Sword, and Colt + Salvo are registered trademarks of Loveland Products, Inc

Buctril and Rimfire are registered trademarks of Bayer CropScience LP

Everest is a registered trademark of Arysta LifeScience North America LLC

BOLT, GoldSky, Starane NXT, and WideMatch are registered trademarks of Corteva Agriscience United States

**SL-4027 070819 07-09-19**

## **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

Notice: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and, to the extent consistent with applicable law, Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

**Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or FMC, and, to the extent permitted by applicable law, Buyer assumes the risk of any such use.**

To the extent consistent with applicable law, FMC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

**This Condition of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.**