



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9629

**FOR DISTRIBUTION AND USE
IN REGISTERED STATES (EXCEPT
NEW YORK AND TEXAS)**

**THIS RECOMMENDATION, WHICH CONTAINS
ADDITIONAL DIRECTIONS FOR USE, IS MADE
AS PERMITTED UNDER FIFRA SECTION 2(ee)
AND HAS NOT BEEN SUBMITTED TO OR
APPROVED BY THE US EPA.**

This recommendation for AVAUNT® eVo insect control is valid until December 31, 2025 or until withdrawn, canceled or suspended.

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS
LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA
REGISTERED LABEL MUST BE FOLLOWED.**

**THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE
APPLICATION.**

FOR SUPPRESSION OF LEAFMINERS AND CONTROL OF LOOPERS IN SUCCULENT BEANS

CROP	APPLICATION METHOD	TARGET PEST	RATE	PHI (Days)	REI (Hours)
Bean, Succulent (except soybean) Including: bean (Phaseolus) (includes lima bean, green; broad bean, succulent; runner bean, snap bean, wax bean); bean (Vigna) (includes asparagus bean, blackeyed pea, Chinese longbean, cowpea, moth bean, southern pea, yardlong bean); jackbean; sword bean	Foliar	Leafminers (suppression only) Loopers	3.5-6.0 oz/A (0.065-0.11 lb ai/A)	3	12



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104

Directions For Use:

- AVAUNT eVo insect control is recommended as a foliar application for suppression of leafminers and control of loopers in succulent beans.
- Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of plants and density of foliage being treated.
- For best performance, use an adjuvant that is approved for use on agricultural crops and follow the directions on the manufacturers label.
- Apply AVAUNT eVo insect control at 3.5-6.0 oz/A (0.065-0.11 lb ai/A) as part of a foliar program in conjunction with other insecticides of a different mode of action that are known to be effective on leafminers and loopers.
- Applications targeting leafminers should begin early in the pest infestation when adult leafminer flies are first observed and before the pest population begins to build to damaging levels.
- Applications to control loopers should target early instar larvae to provide best control and limit succulent bean defoliation.
- Use the lower end of the rate range for low or building insect populations and the higher end of the rate range for greater populations and/or dense foliage.
- Make repeat applications on a 7-day spray schedule as necessary based on insect monitoring to maintain control.
- See the Resistance Management section of the Avaunt eVo insect control Section 3 Federal label for additional guidance regarding insecticide resistance management.

RESTRICTIONS AND PRECAUTIONS

- Do not apply more than 24 oz/A AVAUNT eVo insect control or 0.44 lb ai of indoxacarb-containing products per acre per crop.
- Do not make more than 4 applications per crop.
- Do not apply more than 72 oz/A AVAUNT eVo insect control or 1.32 lb ai of indoxacarb-containing products per acre per year.
- For ground applications, make a uniform application in approximately 20-100 gallons of water per acre.
- For aerial applications, make a uniform application in a minimum of 5 gallons of water per acre.
- The minimum interval between sprays is 7 days.

R-4466 071620 07-13-20

FMC, the FMC logo and Avaunt are trademarks of FMC Corporation or an affiliate
©2020 FMC Corporation



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104