



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9629

**FOR DISTRIBUTION AND USE
ONLY IN THE FOLLOWING
STATES: CALIFORNIA and ARIZONA**

**THIS RECOMMENDATION, WHICH CONTAINS
ADDITIONAL DIRECTIONS FOR USE, IS MADE AS
PERMITTED UNDER FIFRA SECTION 2(ee) AND
HAS NOT BEEN SUBMITTED TO OR APPROVED
BY THE US EPA.**

This recommendation for AVAUNT® EVO INSECT CONTROL is valid until January 15, 2029, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED SUBJECT TO THE TERMS AND CONDITIONS OF THE PRODUCT LABELING.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR THE SUPPRESSION OF LEAF FOOTED PLANT BUGS, GREEN STINK BUGS, *Carpophilus truncatus*, AND FEEDING DAMAGE PROTECTION IN NUTMEAT IN TREE NUTS

CROP	PEST	RATE	APPLICATION DIRECTIONS	REI
Tree Nuts Including: Almond Pistachio cultivars, varieties, and/or hybrids of these	Leaf footed plant bugs, green Stink bugs, and <i>Carpophilus truncatus</i>	6.0 oz/a (0.11 lb a.i.)	Apply as required by scouting, usually from late dormancy until Mid-May. Timing and frequency of applications should be based on insect population and threshold.	12 hrs.

TREE NUTS RESTRICTIONS:

- Make no more than 3 applications per season.
- Do not apply more than 18 oz/A of Avaunt® eVo insect control or 0.33 lb ai/A of indoxacarb-containing products per calendar year.
- Do not apply Avaunt® eVo insect control later than May 15 due to MRLs constrain with EU. It is the grower and processor responsibility to check for residue before shipping overseas.



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000

- The minimum interval between treatments is 7 days.
- Spray Volume: Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of trees or plants and density of foliage.
- Ground Application: Apply in a minimum of 50 gal/A of water by conventional ground equipment. For best results, apply 100-150 gal/A of water.
- Aerial Application: Apply in a minimum of 10 gal/A of water. Applying by air can result in reduced canopy coverage, which may also reduce efficacy. For best results, use MSO or COC blend adjuvants to help increase coverage, penetration and thus performance.
- This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging in the treatment area.

R-4767 010225 010225



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000