



| | | | |
|------------|-------|-----|-------------|
| INDOXACARB | GROUP | 22A | INSECTICIDE |
|------------|-------|-----|-------------|

EPA Reg. No. 279-9629

SUPPLEMENTAL LABELING

FOR FOLIAR APPLICATIONS TO COFFEE AND STRAWBERRY TO CONTROL LISTED PESTS

**THIS SUPPLEMENTAL LABEL EXPIRES 08/31/2027 AND MUST NOT BE USED OR
DISTRIBUTED AFTER THIS DATE**

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read the label affixed to the container for Avaunt® eVo insect control before applying.

Use of Avaunt® eVo insect control according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Avaunt® eVo insect control.

IMPORTANT

**BEFORE USING AVAUNT® eVo insect control, READ AND FOLLOW ALL APPLICABLE DIRECTIONS;
RESTRICTIONS; AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.**

This bulletin contains new or supplemental instructions for use of this product which does not appear on the package label. Follow the instructions carefully.

This label must be in the possession of the user at the time of pesticide application.

Sold By



**FMC Corporation
2929 Walnut Street
Philadelphia, PA 1910**

| Crops | Insects | AVAUNT® eVo insect control Rate Per Acre | | Last Application (Days to Harvest) | REI |
|--------|--------------------|------------------------------------------|--------|---------------------------------------|---------|
| | | lb ai | Ounces | | |
| Coffee | Coffee berry borer | 0.11 | 6.0 | 14 | 12 hrs. |

COFFEE RESTRICTIONS

Make no more than 4 applications per acre per crop.

Do not apply more than 24 oz/A of AVAUNT® eVo insect control or 0.45 lb ai/A of indoxacarb-containing products per calendar year.

The minimum interval between sprays is 7 days.

Note: A commercially appropriate and locally available adjuvant may be added to the tank mixture at a rate listed on the label.

Time applications early in the pest infestation when no more than 2% of the coffee berries are infested with coffee berry borer in position A or B (prior to borer reaching the endosperm/seed). Calibrate equipment to achieve thorough spray coverage of the berry without runoff.

| Crops | Insects | AVAUNT® eVo insect control Rate Per Acre | | Last Application (Days to Harvest) | REI |
|------------|-------------------------------------------------|------------------------------------------|-----------|---------------------------------------|---------|
| | | lb ai | Ounces | | |
| Strawberry | Beet armyworm Cabbage looper Corn earworm | 0.065 - 0.11 | 3.5 - 6.0 | 1 | 12 hrs. |
| | Lygus spp.* | 0.09 – 0.11 | 5.0 – 6.0 | | |

STRAWBERRY RESTRICTIONS

Make no more than 4 applications per acre per crop.

Do not apply more than 24 oz/A of AVAUNT® eVo insect control or 0.45 lb ai/A of indoxacarb-containing products per calendar year.

The minimum interval between sprays is 7 days.

Note: Not all varieties of strawberries have been tested for crop safety with AVAUNT® eVo Insect Control alone or in tank mixture, see "Tank Mixtures and Crop Safety" section for more information.

*Suppression only

(Derived from AVAUNT® eVo insect control label SL-4661A 082124 08-22-24)

R-4710 082124 08-22-24