



FIFRA 2(EE) Recommendation
EPA Reg. No. 279-9638
FOR DISTRIBUTION AND USE ONLY IN REGISTERED STATES (EXCEPT TEXAS)
THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(EE) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE U.S. EPA.

This recommendation is valid until December 31, 2030 or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.
THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

For Suppression of Anthracnose Stalk Rot, Fusarium Stalk and Crown Rot and Physoderma Brown Spot in Corn (Field, Corn Grown for Seed, Popcorn)

Directions For Use

Apply 0.68 fl. oz./1000 row-ft. Xyway™ 3D fungicide (11.8 fl. oz./A on 30" rows) at-planting, in-furrow with the 3rive 3D application system for suppression of anthracnose stalk rot, stalk and crown rot diseases caused by susceptible *Fusarium* spp., and Physoderma brown spot.

CROP	PEST	RATE OF APPLICATION		
		Fluid OZ/Acre (30" Row Spacing)	Fluid OZ/1000 Linear Ft.	Lbs. ai/A
CORN (Field, Corn Grown for Seed, Popcorn)	Anthracnose Stalk Rot (<i>Colletotrichum graminicola</i>)	11.8	0.68	0.228
	Fusarium Stalk and Crown Rot (<i>Fusarium graminearum, Fusarium spp.</i>)			
	Physoderma Brown Spot (<i>Physoderma maydis</i>)			
Restrictions: <ul style="list-style-type: none"> • DO NOT apply more than 0.228 lb. ai/A (11.8 fl. oz. of product/A) per year including at-plant plus foliar applications of other flutriafol-containing products. • DO NOT apply more than one application per year. • DO NOT apply this product as a foliar application. 				

R-5022 012826 01-28-26 (replaces R-4519 012621 10-3-19)
Xyway and FMC – Trademarks of FMC or an affiliate



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000