

## RESTRICTED USE PESTICIDE

Due to toxicity to fish and aquatic organisms For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.



### FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9652

**FOR DISTRIBUTION AND USE  
ONLY IN COLORADO, IDAHO,  
NEBRASKA, OREGON,  
WASHINGTON STATE, AND  
WYOMING**

**THIS RECOMMENDATION, WHICH CONTAINS  
ADDITIONAL DIRECTIONS FOR USE, IS  
MADE AS PERMITTED UNDER FIFRA  
SECTION 2(ee) AND HAS NOT BEEN  
SUBMITTED TO OR APPROVED BY THE US  
EPA.**

This recommendation for ELEVEST™ insect control is valid until December 31, 2026, or until withdrawn, canceled or suspended.

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED. SUBJECT TO THE TERMS AND CONDITIONS OF THE PRODUCT LABELING.**

**THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**

### **FOR CONTROL OF COLORADO POTATO BEETLE ON VOLUNTEER POTATOES AND OTHER WEEDS IN DRY/SUCCULENT BEANS AND PEAS**

#### **Directions For Use**

##### **Pests Controlled, Application Rates, and Application Method**

ELEVEST insect control is recommended for control of Colorado potato beetle on volunteer potatoes, or other weeds, in dry/succulent beans and peas. Apply ELEVEST insect control at 5.6 - 9.6 fl. oz./A (0.098 – 0.167 lb ai/A) as a foliar spray.

##### **Application Timing and Resistance Management**

Apply just prior to or just after egg hatch while larvae are small. In some areas, where local populations of Colorado potato beetle have elevated levels of resistance to insecticides, use ELEVEST insect control at the 9.6 fl. oz/A application rate.

Do not apply ELEVEST insect control more than twice to a generation of Colorado potato beetle or within any 30 day period. Application(s) to the next generation of Colorado potato beetle must be made with an effective product with a different mode of action. With resistant populations of Colorado potato beetle, back-to-back applications on a 5 to 7-day interval may be required to achieve maximum control. See the RESISTANCE MANAGEMENT section of the ELEVEST insect control Section 3 Federal label for additional guidance on insecticide resistance management.

#### **Use Restrictions**

For aerial application, use a minimum of 5 gallons per acre (gpa) of water.

Make no more than 2 applications per acre per calendar year.

Minimum interval between treatments is 3 days for succulent and 7 days for dried beans and peas.

PHI is 3 days for succulent and 14 days for dried beans and peas.

Do not apply more than 19.2 fl oz ELEVEST insect control or 0.2 lb ai. of chlorantraniliprole and 0.2 lb ai of bifenthrin containing products per acre per calendar year.

When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray.

R-4593 061721 06-17-21



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104  
215-299-6000