

PRINCIPAL DISPLAY PANEL

GROUP	2	HERBICIDE
GROUP	4	HERBICIDE

DB-8454 HERBICIDE

Wettable Granules

FOR SALE FOR USE ON WHEAT (SPRING AND DURUM) AND SPRING BARLEY IN THE
PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE
RIVER REGION) ONLY

COMMERCIAL (AGRICULTURAL)

REGISTRATION NO. 28894 PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENTS: Dicamba (present as sodium salt) 53.8%
Thifensulfuron methyl 7.7%
Tribenuron methyl 3.9%

Warning, contains the allergens milk and sulfites



CAUTION

POISON

EYE IRRITANT

NET CONTENTS: 0.2 – 600.0 Kg

READ THE LABEL AND ACCOMPANYING BOOKLET BEFORE USING

FMC of Canada Limited
6755 Mississauga Road, Suite 204
Mississauga, ON L5N 7Y2
1-833-362-7722

PRECAUTIONS:

- KEEP OUT OF REACH OF CHILDREN.
- AVOID BREATHING SPRAY MIST. USE WITH ADEQUATE VENTILATION.
- AVOID CONTACT WITH SKIN, EYES AND CLOTHING. WEAR GOGGLES OR FACE SHIELD, COVERALLS, AND SUITABLE CHEMICAL RESISTANT GLOVES DURING MIXING AND LOADING.
- MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. CAUSES EYE AND SKIN IRRITATION.
- WHEN USING, DO NOT EAT, DRINK OR SMOKE. WASH SKIN THOROUGHLY WITH SOAP AND WATER AFTER HANDLING.
- REMOVE CONTAMINATED CLOTHING IMMEDIATELY AFTER USE. STORE AND WASH CONTAMINATED CLOTHING SEPARATELY FROM HOUSEHOLD LAUNDRY BEFORE REUSE.
- WEAR LONG-SLEEVED SHIRT, LONG PANTS, SOCKS, AND CHEMICAL-RESISTANT GLOVES AND FOOTWEAR DURING MIXING, LOADING, APPLICATION, CLEAN-UP AND REPAIR. CHEMICAL-RESISTANT GLOVES ARE NOT REQUIRED WHILE OPERATING GROUNDBOOM SPRAYERS.
- DO NOT ENTER OR ALLOW WORKER ENTRY INTO TREATED AREAS DURING THE RESTRICTED-ENTRY INTERVAL (REI) OF 12 HOURS.
- DO NOT USE HUMAN FLAGGERS.

FIRST AID:

In case of poisoning, contact a physician or a poison control centre.

IF IN EYES:

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IN CASE OF EMERGENCY, CALL TOLL FREE, DAY OR NIGHT: 1-800-331-3148

TOXICOLOGICAL INFORMATION:

Dicamba may cause severe irritation to the eyes, and irritation to the skin and mucous membranes. Symptoms of overexposure to dicamba may include dizziness, muscle weakness, loss of appetite, weight loss, vomiting, decreased heart rate, shortness of breath, excitement, tenseness, depression, incontinence, cyanosis, muscle spasms, exhaustion and loss of voice. Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS:

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under **DIRECTIONS FOR USE**. Refer to attached booklet for more information.

STORAGE:

Store this product away from food or feed. Store product in original container only, away from other pesticides or fertilizer. Not for use or storage in or around the home. Keep container closed.

DISPOSAL:

If this is a refillable container:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

If this is a returnable container:

DO NOT reuse this container for any other purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Directions For Use Booklet

GROUP	2	HERBICIDE
GROUP	4	HERBICIDE

DB-8454 HERBICIDE

Wettable Granules

FOR SALE FOR USE ON WHEAT (SPRING AND DURUM) AND SPRING BARLEY IN THE
PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE
RIVER REGION) ONLY

COMMERCIAL (AGRICULTURAL)

REGISTRATION NO. 28894 PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENTS: Dicamba (present as sodium salt) 53.8%
Thifensulfuron methyl 7.7%
Tribenuron methyl 3.9%

Warning, contains the allergens milk and sulfites



CAUTION

POISON

EYE IRRITANT

READ THE LABEL AND THIS ACCOMPANYING BOOKLET BEFORE USING

FMC of Canada Limited
6755 Mississauga Road, Suite 204
Mississauga, ON L5N 7Y2
1-833-362-7722

PRECAUTIONS:

- KEEP OUT OF REACH OF CHILDREN.
- AVOID BREATHING SPRAY MIST. USE WITH ADEQUATE VENTILATION.
- AVOID CONTACT WITH SKIN, EYES AND CLOTHING. WEAR GOGGLES OR FACE SHIELD, COVERALLS, AND SUITABLE CHEMICAL RESISTANT GLOVES DURING MIXING AND LOADING.
- MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. CAUSES EYE AND SKIN IRRITATION.

- WHEN USING, DO NOT EAT, DRINK OR SMOKE. WASH SKIN THOROUGHLY WITH SOAP AND WATER AFTER HANDLING.
- REMOVE CONTAMINATED CLOTHING IMMEDIATELY AFTER USE. STORE AND WASH CONTAMINATED CLOTHING SEPARATELY FROM HOUSEHOLD LAUNDRY BEFORE REUSE.
- WEAR LONG-SLEEVED SHIRT, LONG PANTS, SOCKS, AND CHEMICAL-RESISTANT GLOVES AND FOOTWEAR DURING MIXING, LOADING, APPLICATION, CLEAN-UP AND REPAIR. CHEMICAL-RESISTANT GLOVES ARE NOT REQUIRED WHILE OPERATING GROUNDBOOM SPRAYERS.
- DO NOT ENTER OR ALLOW WORKER ENTRY INTO TREATED AREAS DURING THE RESTRICTED-ENTRY INTERVAL (REI) OF 12 HOURS.
- DO NOT USE HUMAN FLAGGERS.

FIRST AID:

In case of poisoning, contact a physician or a poison control centre.

IF IN EYES:

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container or label or the product name and Pest Control Product Registration Number with you when seeking medical attention.

IN CASE OF EMERGENCY, CALL TOLL FREE, DAY OR NIGHT: 1-800-331-3148

TOXICOLOGICAL INFORMATION:

Dicamba may cause severe irritation to the eyes, and irritation to the skin and mucous membranes. Symptoms of overexposure to dicamba may include dizziness, muscle weakness, loss of appetite, weight loss, vomiting, decreased heart rate, shortness of breath, excitement, tenseness, depression, incontinence, cyanosis, muscle spasms, exhaustion and loss of voice. Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS:

- **TOXIC** to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.
- This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

SPRAY DRIFT PRECAUTIONS

DB-8454 Herbicide may cause injury to desirable trees and plants, particularly flowers, fruit trees, grapes, ornamentals, peas, potatoes, tomatoes, and other broadleaf plants, especially in their developmental and growing stage. Follow these precautions when spraying in the vicinity of sensitive crops:

1. Avoid spraying when winds are towards sensitive crops.
2. Use coarse sprays since they are less likely to drift than fine sprays. Select nozzles which minimize amounts of fine spray particles. Keep spray pressure below 150 kPa and the spray volume above 220 L/ha unless otherwise required by the nozzle manufacturer.
3. Do not spray when the temperature is expected to exceed 30°C.
4. Avoid spraying under conditions of high humidity or fog.
- 5

TABLE OF CONTENTS:

PRINCIPAL DISPLAY PANEL	1
DB-8454 HERBICIDE	1
PRECAUTIONS:	2
FIRST AID:	2
IF IN EYES:	2
IF ON SKIN OR CLOTHING:	2
IF SWALLOWED:	2
IF INHALED:	2
TOXICOLOGICAL INFORMATION:	3
STORAGE:	3
DISPOSAL:	3
NOTICE TO USER:	3
DIRECTIONS FOR USE BOOKLET	3
PRECAUTIONS:	4
FIRST AID:	5
IF IN EYES:	5
IF ON SKIN OR COTHING:	5
IF SWALLOWED:	5
IF INHALED:	5
ENVIRONMENTAL PRECAUTIONS:	6
SPRAY DRIFT PRECAUTIONS	6

PRODUCT DESCRIPTION:	7
REPLANTING TO OTHER CROPS:	8
GRAZING/HARVEST RESTRICTIONS:	8
VARIETY RESTRICTIONS:	8
RAINFESTNESS INFORMATION:	8
APPLICATION INFORMATION:	8
APPLICATION PRECAUTIONS:	8
DIRECTIONS FOR USE:	9
APPLICATION INSTRUCTIONS:	9
AERIAL APPLICATION:	ERROR! BOOKMARK NOT DEFINED.
SPRAY BUFFER ZONES:	11 WHEAT (SPRING AND DURUM) AND SPRING BARLEY
	11
DB-8454 Herbicide	12
WHEAT (SPRING AND DURUM) ONLY	13
DB-8454 Herbicide + HORIZON* NG Herbicide	13
MIXING INFORMATION:	14
MIXING PRECAUTIONS:	14
MIXING DIRECTIONS:	14
SPECIAL MIXING INSTRUCTIONS FOR CHEMICAL HANDLERS	14
EQUIPMENT CLEANING INFORMATION:	15
CLEANING PRECAUTIONS:	15
CLEANING INSTRUCTIONS:	15
RESISTANCE-MANAGEMENT RECOMMENDATIONS:	16
STORAGE:	16
DISPOSAL:	16
NOTICE TO USER:	17

PRODUCT DESCRIPTION:

DB-8454 Herbicide is recommended for selective post-emergent control or suppression of broadleaf weeds in wheat (spring and durum) and spring barley, not underseeded to legumes or grasses.

DB-8454 Herbicide contains three active ingredients formulated as dry flowable or water soluble granules. It is non-corrosive, non-flammable and does not freeze. Should product be exposed to freezing temperatures, warm to room temperature prior to use.

DB-8454 Herbicide is to be mixed with water and a recommended adjuvant and applied as a uniform broadcast spray. Recommended adjuvants are: AGRAL90*, AG-SURF* and CITOWETT* PLUS. Apply early postemergence, to the main flush of actively growing broadleaf weeds. APPLY WITH GROUND EQUIPMENT ONLY.

Warm, moist growing conditions promote active weed growth and enhance the activity of DB-8454 Herbicide by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and regrowth may occur. For best results, ensure thorough spray coverage of target weeds. See "DIRECTIONS FOR USE" section of this label for complete use details.

DB-8454 Herbicide stops growth of susceptible weeds rapidly. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

IMPORTANT INFORMATION: READ BEFORE USING

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

- Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, recreational grounds or similar areas.
- Prevent drift of spray to desirable plants.
- Carefully observe sprayer clean-up instructions, as spray tank residues may damage crops other than spring wheat and spring barley.

REPLANTING TO OTHER CROPS:

Any crop may be planted in the year following the use of DB-8454 Herbicide.

GRAZING/HARVEST RESTRICTIONS:

Wheat (spring and durum) and spring barley may not be grazed or fed to livestock within 7 days of application.

VARIETY RESTRICTIONS:

Limit use of DB-8454 Herbicide to currently licensed varieties of wheat (spring and durum) and spring barley. Consult FMC prior to use on unlicensed varieties for crop tolerance information. DO NOT USE ON LEGER SPRING BARLEY OR BELVEDERE WHEAT VARIETIES.

RAINFALL INFORMATION:

Up to 25 mm of rain beginning 4 hours or more after spraying will not reduce the effectiveness of DB-8454 Herbicide.

APPLICATION INFORMATION:

APPLICATION PRECAUTIONS:

- Do not apply to irrigated land where tail water will be used to irrigate cropland.
- Apply DB-8454 Herbicide when air temperature is between 10 and 25°C. Do not apply when there is a risk of severe drop in night temperature.
- Do not treat areas where soil penetration or surface washing may bring DB-8454 Herbicide into contact with roots of desirable plants.
- Do not remove soil from treated fields for use on lawns, in gardens or for back fill.
- Crop damage may occur if DB-8454 Herbicide is applied at any time other than the recommended crop stage.
- Apply with an adjuvant registered for use with DB-8454 Herbicide. Use only water as a carrier. Other carriers may accelerate breakdown of DB-8454 Herbicide and reduce its effectiveness.
- Do not apply to crops underseeded to legumes or grasses as injury to forage may result.
- Application to crops that are stressed by severe weather conditions, frost, low fertility, water logged soil (soil at or near field capacity), disease or insect damage may result in crop injury. Drought, disease or insect damage following application may also result in crop injury.
- Under certain conditions, such as heavy rainfall, prolonged cool weather, frost or wide fluctuations

in day/night temperatures, temporary lightening in crop colour, and occasionally, a slight reduction in crop height may occur.

- Do not apply more than 130 grams/hectare of DB-8454 Herbicide per year.
- Apply only to agricultural crops when the potential for drift to areas of human habitation and areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature, application equipment and sprayer settings.

Note: Crops growing under stress from adverse environmental conditions such as excess moisture, drought, disease, etc. may suffer a further setback and exhibit more pronounced injury symptoms if DB-8454 Herbicide is applied. However, the crop injury that may occur is usually offset by the weed control obtained.

DIRECTIONS FOR USE:

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

APPLICATION INSTRUCTIONS:

Apply the spray mixture uniformly with properly calibrated ground equipment only. Ensure thorough coverage and a uniform spray pattern. Flat fan nozzles are recommended (minimum 55 litres spray volume per hectare). Use 50 mesh or larger filter screens (metal or nylon). A spray pressure of 210-275 kPa is recommended. Avoid overlapping and shut off spray boom while starting, turning, slowing or stopping to prevent crop injury from over application.

Field Sprayer Application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium/coarse classification. Boom height must be 60 cm or less above the crop or ground.

Aerial Application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

Apply only by fixed-wing or rotary aircraft equipment, which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Use Precautions:

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Operator Precautions:

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted. It is desirable that the pilot have communication capabilities at each treatment site at the time of application. The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label. All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions:

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-833-362-7722 or obtain technical advice from the distributor or your provincial agricultural representative.

Application of this specific product must meet and/or conform to the following:

Apply a minimum of 25 L of spray volume per hectare.

Apply a maximum of 50 L of spray volume per hectare.

Spray Buffer Zones

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label,
- low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage,
- soil drench or soil incorporation.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Spray Buffer Zones (metres) Required for the Protection of:		
			Aquatic Habitat of Depths:		Terrestrial Habitat:
			Less than 1 m	Greater than 1 m	
Field sprayer	Wheat (Spring, Durum) and Spring Barley		1	0	15
Aerial	Wheat (Spring, Durum) and Spring Barley	Fixed wing	1	0	125
		Rotary wing	1	0	100

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

TANK MIXES:

This product may be tank mixed with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact FMC of Canada Limited at 1-833-362-7722 for information before applying any tank mix that is not specifically recommended on this label.

WHEAT (SPRING AND DURUM) and SPRING BARLEY

DB-8454 Herbicide	
LOCATION: For use in the Prairie provinces and Interior of British Columbia (including the Peace River region) only.	
CROP(S): Wheat (spring and durum) and spring barley only, not underseeded to legumes.	
TIMING: Post-emergent application from the 2-5 leaf stage of spring wheat or spring barley; seedling stage to 8 leaf stage of kochia.	
TANK MIX COMPONENTS AND APPLICATION RATES	
DB-8454 Herbicide	130 grams/hectare
Adjuvants: AGRAL [®] 90, AG-SURF [®] or CITOWETT [®] PLUS	2 litres/1000 litres of spray solution
MIXING: Ensure that DB-8454 Herbicide is completely dissolved, then add adjuvant. Refer to MIXING DIRECTIONS section for general instructions. Use minimum volume of 55 L/ha for ground and 25 to 50 L/ha for aerial application.	
WEEDS CONTROLLED	
Weeds Controlled	Stinkweed
Annual smartweed (green smartweed, lady's thumb)	Tartary buckwheat
Ball mustard	Volunteer rapeseed ²
Chickweed ¹	Volunteer sunflower
Corn spurry	Wild buckwheat ³
Cow cockle	Wild mustard
Dandelion (spring or fall rosettes, less than 15 cm in diameter)	
Flixweed	Weeds Suppressed
Groundsel (common)	Canada thistle ⁴
Hemp-nettle	Cleavers (1-3 whorls)
Kochia (including Group 2 resistant kochia).	Round leaved mallow ⁵
Lamb's-quarters	Scentless chamomile
Narrow-leaved hawk's beard	Sow thistle ⁴
Redroot pigweed	Stork's bill (2-6 leaf)
Russian thistle	Toadflax ⁶
Shepherd's purse	
NOTES: Do not permit lactating dairy animals to graze fields within 7 days after application. DO NOT harvest forage or cut hay within 30 days after application. Withdraw meat animals from treated fields at least 3 days before slaughter.	

Notes:

1. Apply when chickweed is small (1-6 leaf) and actively growing, but before crop canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.

2. DB-8454 Herbicide alone will not control imazethapyr tolerant canola varieties (e.g. Clearfield* varieties or other varieties with the Pursuit Smart* trait)
3. Apply to actively growing wild buckwheat in the cotyledon to 5-leaf stage. Under environmental stress, such as dry weather, control may be reduced. Large plants may continue to grow after treatment.
4. Apply when the majority of thistles have emerged and are actively growing. For best top growth control, apply before bud stage when thistles are no larger than 15 cm in height and before crop canopy prevents thorough coverage of weeds. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be controlled.
5. Apply to actively growing round leaved mallow from the 2 - 6 leaf stage (10-12 cm in height). DB-8454 Herbicide will keep mallow stunted, but may not reduce overall populations.
6. Apply to toadflax that is no larger than 15 cm in height. A control program for this weed includes both frequent tillage and chemical application.

WHEAT (SPRING AND DURUM) ONLY

DB-8454 Herbicide + HORIZON* NG Herbicide	
LOCATION: For use in the Prairie provinces and Interior of British Columbia (including the Peace River region) only.	
CROP: Wheat (spring and durum) only, not underseeded to legumes	
TIMING: Post-emergent application from the 2-5 leaf stage of spring wheat; seedling stage to 8-leaf stage of kochia.	
TANK MIX COMPONENTS AND APPLICATION RATES	
DB-8454 Herbicide	130 grams/hectare
HORIZON* NG Herbicide	0.930 L/Ha
MIXING: Ensure that DB-8454 Herbicide is completely dissolved. Reduce agitation, and then add HORIZON* NG Herbicide. Refer to MIXING DIRECTIONS section for general instructions. Use minimum volume of 55 L/ha for ground and 30 L/ha for aerial application.	
WEEDS CONTROLLED	
Ground: Broadleaf weeds controlled or suppressed by DB-8454 Herbicide alone, green foxtail and wild oats. Aerial: Broadleaf weeds controlled or suppressed by DB-8454 Herbicide alone, as well as wild oats	
NOTES: Do not permit lactating dairy animals to graze fields within 7 days after application. DO NOT harvest forage or cut hay within 30 days after application. Withdraw meat animals from treated fields at least 3 days before slaughter.	

MIXING INFORMATION:

MIXING PRECAUTIONS:

Do not allow the spray mixture to remain in the tank for more than 24 hours before spraying or the effectiveness may be reduced. If the spray mixture has been allowed to stand in the tank, use vigorous agitation to thoroughly disperse before resuming spraying.

MIXING DIRECTIONS:

Select a spray volume that will ensure thorough coverage and a uniform spray pattern. Use a minimum of 55 L of spray volume per hectare.

1. Always start with a clean and empty sprayer tank.
2. Fill the tank to 2/3 full of **clean** water.
3. With the agitator running, add the required amount of DB-8454 Herbicide. Continue to agitate for a minimum of 5 minutes to ensure that DB-8454 Herbicide is **completely** dissolved.
4. Make sure there are no undissolved granules, and then add the required amount of the tank mix partner. If the tank mix partner is an Emulsifiable Concentrate (EC), reduce agitation to avoid inducing an invert emulsion.
5. Add the rest of the water.
6. If required for the tank mixture, add the appropriate adjuvant. If an antifoam agent is required, add last.
7. Refer to the tank mix sections of this booklet for mixing order and other mixing instructions.
8. For repeat tank loads, reduce the material remaining in the tank to 10% of the original volume **or less** before proceeding with step 1, because remaining chemicals may prevent DB-8454 Herbicide granules from completely dissolving or dispersing. If this is not possible, pre-slurry DB-8454 Herbicide in a small amount (5-10 L) of water before adding to the tank.

SPECIAL MIXING INSTRUCTIONS FOR CHEMICAL HANDLERS

1. Ensure the chemical handler is clean (rinsed with ammonia).
2. For best results, add DB-8454 Herbicide to the top of the sprayer tank.

3. If a chemical handler is used, make sure that all of the granules of DB-8454 Herbicide are completely dissolved/dispersed and injected into the main tank with agitation before adding other chemicals.
4. Low water volumes in some chemical handlers can cause the water to become saturated with DB-8454 Herbicide, leaving undissolved granules. In that case, rinse the chemical handler with clean water before adding any other chemical.
5. Always dissolve the DB-8454 Herbicide granules with agitation in the main tank before adding a tank mix partner.

NOTE: Follow the clean-out procedure recommended for sprayer tanks for the chemical handler.

EQUIPMENT CLEANING INFORMATION:

CLEANING PRECAUTIONS:

CAUTION: Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour, which may cause eye, nose, throat, and lung irritation. Do not clean equipment in an enclosed area.

DO NOT CLEAN SPRAYER NEAR WELL OR WATER SOURCE OR NEAR TO DESIRABLE VEGETATION.

CLEANING INSTRUCTIONS:

To avoid subsequent injury to crops other than wheat (spring and durum) or spring barley, immediately after spraying and prior to spraying other crops, thoroughly remove all traces of DB-8454 Herbicide from mixing and spray equipment as follows:

1. Drain tank and then fill tank half-full with clean water. Flush tank, boom and hoses by operating the sprayer until the system is purged of the rinse water. If boom is equipped with multiple nozzle bodies, be sure to rotate through all nozzles to ensure clean water reaches all parts of these assemblies.
2. Visually inspect tank to ensure removal of all visible herbicide residues. If necessary, repeat Step 1.
3. Fill the tank with clean water, and then add 1 litre household AMMONIA (containing minimum of 3% ammonia) or equivalent amount of sprayer tank cleaner containing ammonia, per 100 litres of water. Ammonia will not neutralize the herbicide, but helps dissolve dried-on herbicide deposits. If an emulsifiable concentrate has been tank mixed with DB-8454 Herbicide, use of a wetting agent in the cleaning process will assist in removing oily residue that may trap DB-8454 Herbicide on the tank and hoses.
4. Flush solution through boom and hoses, and then add more water to completely fill tank. Allow to sit for 15 minutes with agitation.
5. Drain the tank.
6. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
7. Flush the sprayer with two tanks full of clean water, flushing water through the hoses and boom.

RESISTANCE-MANAGEMENT RECOMMENDATIONS:

For resistance management, DB-8454 Herbicide is a Group 2 and Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to DB-8454 Herbicide and other Group 2 and/or Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- ◆ Where possible rotate the use of DB-8454 Herbicide or other Group 2 and Group 4 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- ◆ Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- ◆ Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- ◆ Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- ◆ Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- ◆ For further information or to report suspected resistance, contact your local FMC representative or the FMC hotline at 1-833-362-7722.

STORAGE:

Store this product away from food or feed. Store product in original container only, away from other pesticides or fertilizer. Not for use or storage in or around the home. Keep container closed

DISPOSAL:

If this is a refillable container:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

If this is a returnable container:

DO NOT reuse this container for any other purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean up of spills.

PMRA Approved Label – M.A.
2023-2397 2024-10-10

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

®/TM Trademark of FMC Corporation or an affiliate.

* All other products mentioned are trademarks of their respective companies.