



## FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9656

**FOR DISTRIBUTION AND USE  
ONLY IN COLORADO, IDAHO,  
NEBRASKA, OREGON,  
WASHINGTON STATE, AND  
WYOMING**

**THIS RECOMMENDATION, WHICH CONTAINS  
ADDITIONAL DIRECTIONS FOR USE, IS  
MADE AS PERMITTED UNDER FIFRA  
SECTION 2(ee) AND HAS NOT BEEN  
SUBMITTED TO OR APPROVED BY THE US  
EPA.**

This recommendation for VANTACOR™ insect control is valid until December 31, 2026, or until withdrawn, canceled or suspended.

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED. SUBJECT TO THE TERMS AND CONDITIONS OF THE PRODUCT LABELING.**

**THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**

### **FOR CONTROL OF COLORADO POTATO BEETLE ON VOLUNTEER POTATOES AND OTHER WEEDS IN DRY/SUCCULENT BEANS AND PEAS**

#### **Directions For Use**

##### **Pests Controlled, Application Rates, and Application Method**

VANTACOR insect control is recommended for control of Colorado potato beetle on volunteer potatoes, or other weeds, in dry/succulent beans and peas. Apply VANTACOR insect control at 1.2 – 2.5 fl. oz./A (0.047 – 0.098 lb ai/A) as a foliar spray.

##### **Application Timing and Resistance Management**

Apply just prior to or just after egg hatch while larvae are small. In some areas, where local populations of Colorado potato beetle have elevated levels of resistance to insecticides, use VANTACOR insect control at the 2.5 fl. oz/A application rate.

Do not apply VANTACOR insect control more than twice to a generation of Colorado potato beetle or within any 30 day period. Application(s) to the next generation of Colorado potato beetle must be made with an effective product with a different mode of action. With resistant populations of Colorado potato beetle, back-to-back applications on a 5 to 7-day interval may be required to achieve maximum control. See the RESISTANCE MANAGEMENT section of

the VANTACOR insect control Section 3 Federal label for additional guidance on insecticide resistance management.

**Use Restrictions**

Do not make more than 4 applications per acre per crop or 12 applications per acre per calendar year.

Minimum interval between treatments is 3 days.

PHI is 1 day.

Do not apply more than 5.1 fl oz VANTACOR insect control or 0.2 lb a.i. of chlorantraniliprole containing products per acre per crop.

Do not apply more than 15.4 fl oz VANTACOR insect control or 0.6 lb a.i. of chlorantraniliprole containing products per acre per calendar year.

R-4597 061821 06-17-21



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104  
215-299-6000