



## FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9656

**FOR DISTRIBUTION AND USE  
ONLY IN THE STATES OF  
ARKANSAS, KENTUCKY,  
LOUISIANA, MISSISSIPPI,  
MISSOURI, OKLAHOMA and  
TENNESSEE**

**THIS RECOMMENDATION, WHICH  
CONTAINS ADDITIONAL DIRECTIONS FOR  
USE, IS MADE AS PERMITTED UNDER FIFRA  
SECTION 2(ee) AND HAS NOT BEEN  
SUBMITTED TO OR APPROVED BY THE US  
EPA.**

This recommendation for VANTACOR™ insect control is valid until December 31, 2026, or until withdrawn, canceled or suspended.

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED. SUBJECT TO THE TERMS AND CONDITIONS OF THE PRODUCT LABELING.**

**THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**

## **REDUCED RATES FOR CONTROL OF FALL ARMYWORM IN BERMUDAGRASS AND OTHER PASTURE GRASSES**

### **Directions For Use**

VANTACOR insect control is recommended for control of fall armyworm (*Spodoptera frugiperda*) in bermudagrass and other pasture grass forage, fodder and hay.

Apply VANTACOR insect control at 0.9 – 1.1 fluid ounces per acre (0.035 – 0.043 lbs ai/acre) as a foliar spray. Adjuvants such as methylated seed oil (MSO) may be added for improved coverage and residual. Apply when populations reach local established treatment thresholds to prevent crop damage.

Do not make more than two sequential applications of VANTACOR insect control before rotating to another registered insecticide having a different mode-of-action.

See the RESISTANCE MANAGEMENT section of the VANTACOR insect control Section 3 Federal label for additional guidance on insecticide resistance management.



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104  
215-299-6000

## Use Restrictions

Do not apply more than 5.1 fl oz of VANTACOR insect control or 0.2 lb ai chlorantraniliprole containing products per acre per calendar year.

Do not make more than 4 applications per acre per calendar year.

Minimum interval between treatments is 7 days.

Preharvest interval is 0 day.

The field re-entry interval is 4 hours.

R-4622 072121 07-20-21

FMC, the FMC logo and PREVATHON are Trademarks of FMC Corporation or an affiliate.

©2021 FMC Corporation



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104  
215-299-6000