



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9656

**FOR DISTRIBUTION AND USE
ONLY IN THE STATE OF TEXAS**

**THIS RECOMMENDATION, WHICH
CONTAINS ADDITIONAL DIRECTIONS FOR
USE, IS MADE AS PERMITTED UNDER FIFRA
SECTION 2(ee) AND HAS NOT BEEN
SUBMITTED TO OR APPROVED BY THE US
EPA.**

This recommendation for VANTACOR™ insect control is valid until December 31, 2024, or until withdrawn, canceled, or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED. SUBJECT TO THE TERMS AND CONDITIONS OF THE PRODUCT LABELING.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

APPLICATIONS FOR CONTROL OF SESAME LEAFROLLER IN SESAME

Directions For Use

USE DIRECTIONS:

- VANTACOR insect control is recommended for control of sesame leafroller in sesame.
- Apply VANTACOR insect control at 1.2 – 1-7 fluid ounces per acre (0.047 – 0.066 lb ai/A) as a foliar spray.
- For best results, make applications to small larvae and before populations exceed treatment thresholds. Follow local university recommendations for treatment thresholds and best timing.
- Thorough spray coverage is essential for best performance. Use a minimum of 10 gallons by ground or 2 gallons by air. Higher spray volume of 15 gal. by ground and 3 gal. by air will improve control.
- VANTACOR insect control may be mixed with approved agricultural adjuvants in situations where spray coverage is difficult to achieve such as closed canopy or dense foliage. An oil-based adjuvant may improve deposition in hot dry weather. Follow directions on adjuvant manufacturer's label.
- Chemigation: Refer to the VANTACOR insect control Section 3 label for specific directions and precautions for application via chemigation.



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000

- See the RESISTANCE MANAGEMENT section of the VANTACOR insect control Section 3 Federal label for additional guidance on insecticide resistance management.

RESTRICTIONS AND PRECAUTIONS:

- Do not apply more than 5.1 fl oz VANTACOR insect control or 0.2 lb ai of chlorantraniliprole containing products whether applications are made to the soil, foliarly or as a seed treatment per acre per calendar year.
- Do not make more than 4 applications per acre per calendar year.
- Minimum interval between applications is 5 days.
- Preharvest interval is 1 day for sesame.
- The field re-entry interval is 4 hours.

R-4595 080321 06-08-21

FMC, the FMC logo and VANTACOR are Trademarks of FMC Corporation or an affiliate.
©2021 FMC Corporation



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000