

FIFRA 2(ee) Recommendation

EPA Reg. No.: 279-9658

FOR DISTRIBUTION AND USE IN ALL REGISTERED STATES (EXCEPT TEXAS)

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation is valid until March 1, 2028 or until withdrawn, canceled or suspended.

Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

For Suppression of Tar Spot in Corn (Field, Corn Grown for Seed, Popcorn)

Directions For Use

Apply 0.87 fl. oz./1000 row-ft. Xyway™ LFR fungicide (15.2 fl oz/A on 30-in rows) at planting for suppression of tar spot.

CROP	PEST	RATE OF APPLICATION		
		fl oz/A (30-in row spacing)	fl oz/1000 row- feet	Ib ai/acre
CORN (Field, Corn Grown for Seed, Popcorn)	Tar Spot (Phyllachora maydis)	15.2	0.87	0.228

Restrictions:

- Do not apply more than 0.228 lb ai (flutriafol) per acre (15.2 fl oz of product/A) per year including at-plant plus foliar applications of other flutriafol-containing products.
- Do not apply this product as a foliar application.

R-4785 021523

