



FIFRA 2(ee) Recommendation

EPA Reg. No.: 279-9658

FOR DISTRIBUTION AND USE IN THE STATE OF TEXAS

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This label is valid until December 31, 2028, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

SIDE DRESS APPLICATION OF XYWAY® LFR® FUNGICIDE IN CORN, GRAIN SORGHUM

Use Directions for Corn (Post Emergence/Side Dress) (Field, Corn Grown for Grain, Seed or Silage, Popcorn, Sweet Corn), Grain Sorghum

Diseases	USE RATES		
	Fl. Oz./Acre	Fl. Oz./1000 Linear ft. *	Pound ai/A
Corn Diseases: Gray Leaf Spot <i>(Cercospora zeae-maydis)</i> Southern Corn Leaf Blight <i>(Bipolaris maydis)</i> Northern Corn Leaf Blight <i>(Exserohilum turcicum)</i> Common Rust <i>(Puccinia sorghi)</i> Grain Sorghum Diseases: Anthracnose Leaf Blight <i>(Colletotrichum graminicola)</i> Gray Leaf Spot <i>(Cercospora sorghi)</i> Corn Disease Suppression: Physoderma Brown Spot <i>(Physoderma maydis)</i> Tar Spot <i>(Phyllachora maydis)</i>	30" row center = 15.2 fl. oz./A 36" row center = 12.0 fl. oz./A 38" row center = 12.6 fl. oz./A (See label for conversions for other row spacings)	0.87	0.228



FMC Corporation
 2929 Walnut St.
 Philadelphia, PA 19104
 215-299-6000

Side Dress Application:

Ensure product is uniformly dispersed in tank prior to application. Apply using a rate controller system with a hydraulic centrifugal pump and a return line providing tank agitation.

*Refer to the table Xyway LFR® fungicide Required Per Acre for Various Row Spacings” in Application and Mixing Instructions for application rates based on various row spacings.

Apply to the soil after corn or grain sorghum emergence through V6 stage corn or V8 stage grain sorghum. Apply at least 0.5 inches away from the plant, up to 15 inches away from the row. Apply as a solid stream or as a concentrated band application with either water or fluid fertilizer as the carrier to either the soil surface or subsurface injection. Rainfall or irrigation is required after application to incorporate surface applications into the soil for root uptake for maximum performance. Apply the full use rate (0.87 oz/1000 ft of row).and prior to V6 corn V8 grain sorghum for best results.

Restrictions:

- **DO NOT** apply more than 0.228 lb. ai flutriafol/A per year to all corn types or grain sorghum, including at-plant plus foliar applications of other flutriafol-containing products.
- No single application may exceed 15.2 fl. oz. of product (0.228 lb. ai)/A.
- The minimum retreatment interval is 7 days.
- **DO NOT** apply more than 1 side dress application per year.
- Restricted Entry Interval (REI): The REI for detasselling field corn and popcorn grown for seed is 5 days. The REI for sweet corn is 3 days. The REI for all other activities is 12 hours. The REI for grain sorghum is 12 hours.
- Preharvest Interval (PHI): **DO NOT** apply less than 0-day for corn forage harvest, and 7-days for stover, silage and grain of field corn, popcorn and sweet corn. **DO NOT apply less than 30 days before harvest of grain sorghum.**

R-4773 121324 12-13-24_r1



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000