



FIFRA 2(EE) Recommendation
EPA Reg. No. 279-9658
FOR DISTRIBUTION AND USE ONLY IN REGISTERED STATES (EXCEPT TEXAS)
THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(EE) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE U.S. EPA.

This recommendation is valid until December 31, 2030 or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

### For Suppression of Anthracnose Stalk Rot, Fusarium Stalk and Crown Rot and Physoderma Brown Spot in Corn (Field, Corn Grown for Seed, Popcorn)

#### Directions For Use

Apply 0.87 fl. oz./1000 row-ft. Xyway™ LFR fungicide (15.2 fl. oz./A on 30" rows) at-planting for suppression of anthracnose stalk rot, stalk and crown rot diseases caused by susceptible *Fusarium* spp., and *Physoderma* brown spot.

CROP	PEST	RATE OF APPLICATION		
		Fluid OZ/Acre (30" Row Spacing)	Fluid OZ/1000 Linear Ft.	Lbs. ai/A
CORN (Field, Corn Grown for Seed, Popcorn)	<b>Anthracnose Stalk Rot</b> ( <i>Colletotrichum graminicola</i> )	15.2	0.87	0.228
	<b>Fusarium Stalk and Crown Rot</b> ( <i>Fusarium graminearum</i> , <i>Fusarium</i> spp.)			
	<b>Physoderma Brown Spot</b> ( <i>Physoderma maydis</i> )			

Restrictions:

- **DO NOT** apply more than 0.228 lb. of flutriafol/A per year including at-plant plus foliar applications of other flutriafol products.
- No single application may exceed 15.2 fl. oz. of product (0.228 lb. ai)/A.
- **DO NOT** apply more than 1 at plant application per year.
- **DO NOT** apply more than 15.2 fl. oz. of product (0.228 lb. ai)/A per year.
- Restricted Entry Interval (REI): The REI for detasselling field corn and popcorn grown for seed is 5 days. The REI for sweet corn is 3 days. The REI for all other activities is 12 hours.
- Preharvest Interval (PHI): Zero (0) day of harvest for forage, and 7 days for stover and grain of field corn and popcorn and sweet corn.
- Do not apply this product as a foliar application.

R-5024 012826 01-28-26 (replaces R-4521 012621 11-04-20)

Xyway and FMC – Trademarks of FMC or an affiliate



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104  
215-299-6000