

FLUTRIAFOL	GROUP	3	FUNGICIDE
------------	-------	----------	-----------



FIFRA 2(EE) Recommendation
EPA Reg. No. 279-9658
FOR DISTRIBUTION AND USE ONLY IN TEXAS
THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(EE) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE U.S. EPA.

This recommendation is valid until December 31, 2028 or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

At-Plant Off the Seed (2x0, 2x2, 2x2x2 dual placement) of Xyway™ LFR® fungicide in Corn

Directions For Use

Apply Xyway™ LFR® fungicide to corn at planting at least ½” away from the seed, or placed off the row (i.e. 2x0, 2x2, 2x2x2 dual placement, or similar placements) within 4-inches of the seed, but avoiding direct contact with the seed.

- Apply the full use rate (15.2 fl. oz. of product/A) for best results.

Restrictions:

- **DO NOT** apply more than 0.228 lb. of flutriafol/A per year including at-plant plus foliar applications of other flutriafol products.
- No single application may exceed 15.2 fl. oz. of product (0.228 lb ai)/A.
- **DO NOT** apply more than 1 at plant application per year.
- **DO NOT** apply more than 15.2 fl. oz. of product (0.228 lb. ai)/A per year.
- Restricted Entry Interval (REI): The REI for detasselling field corn and popcorn grown for seed is 5 days. The REI for sweet corn is 3 days. The REI for all other activities is 12 hours.
- Preharvest Interval (PHI): Zero (0) day of harvest for forage, and 7 days for stover and grain of field corn and popcorn and sweet corn.

R-5023 012826 01-28-26 (replaces R-4685 013122)

Xyway and FMC – Trademarks of FMC or an affiliate



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000