



Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE IN MINNESOTA, MONTANA, NORTH DAKOTA, AND SOUTH DAKOTA

Preview 2.1 SC Herbicide EPA Reg. No. 70506-394

Fallow or Eco-Fallow Cropland use directions

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by US EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read and follow all applicable instructions, restrictions, and precautions on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

FALLOW OR ECO-FALLOW CROPLAND

Apply Preview 2.1 SC Herbicide in the spring or fall to stubble of harvested crops, or fallow ground for control of weeds prior to emergence in no-till or conservation tillage (CRP) systems. Make application by ground or air, using a minimum of 10 gal/A by ground or 5 gal/A by air.

For fall applications, apply when soil temperatures are 55F or below at a 4-inch soil depth. Apply after September 30 in those areas North of Interstate 90 and after October 15 in those areas North of Interstate 40.

Preview 2.1 SC Herbicide may be tank mixed with other herbicides labeled for this use. The addition of atrazine at 0.5-1.0 lb/A can improve residual control of some weed species. Tank mix a labeled burndown herbicide for control of emerged broadleaf or grass weeds.

<u>USE RATE TABLE FOR FALLOW or ECO-FALLOW CROPLAND</u>		
<u>Spring or Fall¹ Applications for Control of Kochia, Russian Thistle, Lambsquarters and other Labelled Weed Species</u>		
	<u>Broadcast Rate - Fl. Oz. per Acre^{2,3}</u>	
	<u>% Organic Matter</u>	
<u>Soil Texture⁴</u>	<u>1.0 - 2.0%</u>	<u>2.0 - 4.0%</u>
<u>Coarse</u>	<u>11 - 14</u>	<u>14 - 21</u>
<u>Medium</u>	<u>14 - 21</u>	<u>18 - 23</u>
<u>Fine</u>	<u>18 - 23</u>	<u>21 - 26</u>
¹ Refer to section for Fall Applications of PREVIEW 2.1 SC ahead of Field Corn for proper rates and application guidelines when Field Corn is to be planted the following spring		
² Use the higher rate for suppression of grasses and sedges.		



Recommendation

³Refer to rotation guidelines of the PREVIEW 2.1 SC label to ensure the selected use rate of PREVIEW 2.1 SC aligns with rotational intervals and restrictions for subsequent cropping plans

⁴Refer to the previous information on soil types under the SOIL CLASSIFICATION CHART.

Use higher rates for soils of pH less than 7.0 and lowest rate listed per soil type and organic matter for pH greater than 7.0.

Restrictions:

- **DO NOT** apply to frozen soils
- **DO NOT** apply more than a total of 26 fl oz/A of Preview 2.1 SC Herbicide (0.23 lb ai sulfentrazone and 0.45 lb ai metribuzin) as a fallow application per 12 month period.
- **DO NOT** apply more than a total of 0.23 lb ai sulfentrazone per year or 0.45 lb ai of metribuzin per 12 month period.
- **DO NOT** apply to soils classified as sand, with less than 1% organic matter.

****Refer to Crop Rotation table below when applying fall fallow ground applications (Table from Preview 2.1 Label)**

CROP ROTATION

Waiting Period After PREVIEW 2.1 HERBICIDE Application			
Anytime	Tomatoes (Transplanted only)	Soybeans	Sugarcane
4 Months	Barley	Wheat	Field Corn ¹ (Fall application only)
10 Months	Rice		
12 Months	Alfalfa Dry Beans Potato	Asparagus Sunflower Sorghum (2)	Peanuts Tobacco Cotton (5)
18 Months	Cotton (3) And all other crops not listed on the label (3)	Sorghum	Sweet Corn (3)
24 Months	Canola (3)	Sugarbeets (3) (4)	
¹ Field Corn includes corn grown for grain, forage or silage, and seed corn. ² Sorghum may be planted after 12 months where PREVIEW 2.1 HERBICIDE was applied at 34 fl.oz./acre or less in the previous cropping season. ³ Crops that have rotational intervals greater than 12 months after a PREVIEW 2.1 HERBICIDE application are the result of crop injury concerns. ⁴ A rotation interval of 24 months is allowed with a successful bioassay. ⁵ Cotton may be planted after 12 months where PREVIEW 2.1 HERBICIDE was applied at rates 28 fl.oz./acre or less and meets the following conditions: • Medium and fine soils • pH <7.2 • Rainfall or irrigation must exceed 15" after application of PREVIEW 2.1 HERBICIDE to rotate to cotton			