



FIFRA 2(ee) Recommendation
EPA Reg. No. 279-3642
FOR DISTRIBUTION AND USE IN ALL REGISTERED STATES (EXCEPT TX):
THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation for Adastrio™ fungicide is valid until January 31, 2028, or until withdrawn, canceled or suspended.

Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.
THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

For tar spot management in corn (Field Corn, Field Corn Grown for Seed, Sweet Corn and Popcorn)

Directions for Use		
Apply when disease first appears. A second application may be made 10-14 days later. When disease pressure is high, use the higher rate and shorter interval. Apply by aerial or ground application or chemigation.		
Crop	Pest	Rate (fl oz/A)
Corn (Field, Field Corn Grown for Seed, Sweet Corn and Popcorn)	Tar Spot (<i>Phyllachora maydis</i>)	7.0-9.0
RESTRICTIONS: <ul style="list-style-type: none"> • Apply only through R4 (early dough stage) growth stage. • No single application may exceed 9 fl. oz. of product/A. • Do not apply more than 2 applications of product per year. • Do not apply more than 18 fl. oz. of product/A per year. • Do not apply more than 0.228 lb ai of flutriafol/A per year. • Do not apply more than 2.0 lb ai of azoxystrobin/A per year. • Do not apply more than 0.27 lb ai of fluindapyr/A per year. • Do not use an adjuvant after the V8 stage and prior to the VT stage of corn. An adjuvant may be used at any other growth stage. • Restricted Entry Interval (REI): The REI for detasseling field corn and popcorn grown for seed and hand harvesting and detasseling sweet corn is 14 days. The REI for all other activities is 12 hours. • Preharvest Interval: The PHI is 7 days of harvest for forage, and 30 days for stover and grain of field corn and popcorn, and 14 days for sweet corn. 		

R-4764 010323