



FIFRA 2(ee) Recommendation
EPA Reg. No. 279-3642
FOR DISTRIBUTION AND USE ONLY IN: Texas
THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

This recommendation for ADASTRIO™ fungicide is valid until 12/31/2027, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

CROP	PEST	RATE OF APPLICATION
Winter Wheat	<ul style="list-style-type: none"> • Leaf rust (<i>Puccinia triticina</i>, <i>P. hordei</i>) • Stripe rust (<i>Puccinia striiformis</i>) • Stagonospora leaf/glume blotch (<i>Parastagonospora nodorum</i>) • Tan spot (<i>Pyrenophora tritici-repentis</i>) • Septoria leaf blotch (<i>Zymoseptoria tritici</i>) • Stem rust (<i>Puccinia graminis</i>) • Net blotch (<i>Drechslera teres</i>) • Powdery mildew (<i>Blumeria graminis</i>) 	<p>3.5 fl. oz./A - greenup followed by</p> <p>3.5 – 9.0 fl. oz./A – flag leaf</p>

APPLICATION INSTRUCTIONS AND RESTRICTIONS:

- Apply in early spring at greenup, from tillering to early joint (Feekes stages 3-7) with a sequential application occurring at flag leaf (Feekes stages 8-10).
- No single application may exceed 9 fl. oz. of product/A.
- Do not apply more than 2 applications per year.
- Do not apply more than 18 fl. oz. of product/A per year.
- Do not apply more than 0.228 lb ai of flutriafol/A per year.
- Do not apply more than 0.27 lb ai of fluindapyr/A per year.
- Do not apply more than 0.40 lb ai of azoxystrobin/A per year.
- Apply in a minimum of 2 GPA (aerial) and 10 GPA (ground) for optimal coverage and efficacy.
- Restricted Entry Interval (REI) is 12 hours.
- Preharvest Interval: PHI is 7 days prior to harvest of forage, 15 days prior to harvest of hay, and 30 days prior to harvest of grain/straw.

It is strongly recommended that to achieve extended crop protection and manage resistance in the presence of susceptible varieties or heavy disease pressure, full labeled rates of Adastrio (5-9 fl. oz./A), Topguard (10-14 fl. oz./A), Topguard EQ (4-7 fl. oz./A) or Lucento (3-5.5 fl. oz./A) fungicides be applied at flag leaf.

Adastrio fungicide may be mixed with adjuvants, insecticides (such as **Mustang® Maxx** insecticide) or herbicides (such as **Finesse®** Cereal and Fallow herbicide or **Harmony®** Extra SG herbicide) to provide insect or weed control and disease management in one application.



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000

R-4914 111124 11-11-24
ADASTRIO, HARMONY, TOPGUARD, LUCENTO, MUSTANG,
FINESSE and FMC – Trademarks of FMC or an affiliate.
©2024 FMC Corporation. All rights reserved.