

AZOXYSTROBIN	Group	11	Fungicide
FLUINDAPYR	Group	7	Fungicide
FLUTRIAFOL	Group	3	Fungicide

Supplemental Labeling



EPA Reg. No. 279-3642

For use on soybean and for soybean as a rotational crop.

THIS SUPPLEMENTAL LABEL EXPIRES ON May 31, 2027 AND MUST NOT BE USED OR DISTRIBUTED AFTER THIS DATE.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read the entire label. This label must be in the possession of the user at the time of application. Read the label affixed to the container for Adastrio™ before applying. Use of Adastrio™ according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the Adastrio™ container.

SUPPLEMENTAL LABELING

Crop Rotation: Fields treated with Adastrio™ may be planted to the following crops and plant back intervals:

Crop	Plant back Interval
Barley; field corn; field corn grown for seed; sweet corn; popcorn; grain sorghum; soybean; tree nuts; triticale; or wheat	Immediately
Alfalfa; cotton; cucurbits; fruiting vegetables; grapes; hops; Brassica leafy vegetables crop group 5; leafy vegetables, except Brassica, crop group 4; peanut; pome fruit; rice; stone fruit; strawberry; sugar beet; tomato	12 months
All other crops	Prohibited

CROP USE DIRECTIONS

Use Directions for Soybean

Not for use in California

Diseases	Product Rate (fl oz/A)	Application Directions
Anthracnose (<i>Colletotrichum truncatum</i>) Brown spot (<i>Septoria glycines</i>) Cercospora leaf blight and purple seed stain(<i>Cercospora kikuchii</i>) Frogeye leaf spot (<i>Cercospora sojina</i>) Powdery mildew (<i>Erysiphe diffusa</i>) Rust (<i>Phakopsora pachyrhizi</i>) Target spot (<i>Corynespora cassiicola</i>) White mold (<i>Sclerotinia sclerotiorum</i>)	7 to 9	Apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development. Use higher rate and shorter interval when disease pressure is severe.
Restrictions: <ul style="list-style-type: none"> • No single application may exceed 9 fl. oz. of product (0.07 lb fluindapyr, 0.105 lb flutriafol, 0.105 lb azoxystrobin)/A. • Minimum retreatment interval (RTI) is 14 days. • DO NOT apply more than 2 applications per year. • DO NOT apply more than 18 fl oz of product/A per year. • DO NOT apply more than 0.488 lb ai of flutriafol/A per year. • DO NOT apply more than 0.224 lb ai of fluindapyr/A per year • DO NOT apply more than 2.0 lb ai of azoxystrobin/A per year. • Restricted Entry Interval (REI) is 12 hours. • DO NOT feed forage or hay to animals or permit animals to graze. • Preharvest Interval (PHI) is 21 days for seed. 		