



FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9661

FOR THE DISTRIBUTION AND USE ONLY IN
THE STATE OF CALIFORNIA

THIS RECOMMENDATION, WHICH CONTAINS
ADDITIONAL DIRECTIONS FOR USE, IS MADE A
PERMITTED UNDER FIFRA SECTION 2(EE) AND HAS NOT
BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation for Coragen® eVo insect control is valid until December 31, 2030, or until withdrawn, cancelled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.
THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION

FOR USE VIA DRIP CHEMIGATION APPLICATION IN GLOBE ARTICHOKE FOR CONTROL OF BEET ARMYWORM AND CABBAGE LOOPER

Directions For Use

PESTS CONTROLLED

Coragen® eVo insect control is recommended for the control of beet armyworm and cabbage looper via drip chemigation in globe artichoke.

Read the chemigation section of the federal label for more information on how to use via drip chemigation.

See the RESISTANCE MANAGEMENT section of the Coragen® eVo insect control Federal label for additional guidance regarding insecticide resistance management.

Crop	Application Method	Target Pest	Lb. AI Per Acre	Fluid Ounce Product Per Acre	Last Application (Days to Harvest)	REI (Hours)
Artichoke, globe	Drip Chemigation	Beet Armyworm Cabbage Looper	0.047 – 0.098	1.2 – 2.5	3	4

USE RESTRICTIONS

Begin drip application when insect populations first appear. Coragen® eVo insect control must be applied uniformly in the root zone or poor performance will result.

Drip tape must be placed directly underneath a single row to ensure Coragen® eVo is applied in the root zone.

During the first half of the crop growing cycle, allow 1-3 days for Coragen® eVo insect control to be translocated into the leaves for pest protection. For applications made in later stages of plant development, translocation of Coragen® eVo insect control into aerial portions of the plant may take up to 7-10 days.

Do not make more than 2 drip chemigation applications of Coragen® eVo insect control per crop.

Minimum application interval between drip chemigation treatment is 14 days.

Apply no more than 5.1 fl. oz. Coragen® eVo insect control or 0.2 lb. a.i. of chlorantraniliprole containing products per acre per calendar year.

R-4988 081925 08-21-25