



## FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9661

**FOR DISTRIBUTION AND USE  
IN ALL REGISTERED STATES  
EXCEPT NY AND TX**

THIS RECOMMENDATION, WHICH  
CONTAINS ADDITIONAL DIRECTIONS FOR  
USE, IS MADE AS PERMITTED UNDER FIFRA  
SECTION 2(ee) AND HAS NOT BEEN  
SUBMITTED TO OR APPROVED BY THE US  
EPA.

This recommendation for Coragen® eVo insect control is valid until December 31, 2030, or until withdrawn, canceled or suspended.

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS  
LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA  
REGISTERED LABEL MUST BE FOLLOWED.**

**THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE  
APPLICATION.**

## **FOR SUPPRESSION OF LESSER CORNSTALK BORER (*Elasmopalpus lignosellus*) IN SUGARCANE**

<b>Directions For Use</b>					
<b>CROP</b>	<b>APPLICATION METHOD</b>	<b>TARGET PEST</b>	<b>RATE</b>	<b>PHI (Days)</b>	<b>REI (Hours)</b>
Sugarcane	Foliar	Lesser Cornstalk Borer ( <i>Elasmopalpus lignosellus</i> )	1.2-1.7 fl oz/A (0.047-0.066 lb ai/A)	14	4

### **USE DIRECTIONS:**

- Coragen® eVo insect control is recommended for suppression of lesser cornstalk borer (*Elasmopalpus lignosellus*) in sugarcane.
- Apply Coragen® eVo insect control at 1.2-1.7 fluid ounces per acre (0.047-0.066 lb ai/A) as a foliar spray for lesser cornstalk borer infestations in sugarcane. Foliar applications should be made in sufficient water volume to achieve thorough coverage of foliage and the soil area around the sugarcane shoots.



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104  
215-299-6000

- Apply higher rates within the listed range for heavier infestations, larger/denser crop canopy or extreme environmental conditions such as rainy weather and high temperatures.
- For best results, applications of Coragen® eVo insect control by conventional foliar application methods should be followed with an overhead irrigation or rainfall event of 0.5-1.0 inches of water within 24-48 hours after application to enhance contact with lesser cornstalk borer larvae.
- Sugarcane fields with a history of lesser cornstalk borer populations should be scouted weekly and Coragen® eVo insect control applications made early when adult moths are active and laying eggs and when larvae are just beginning to hatch.
- Make applications when populations reach local established treatment thresholds to prevent crop damage.
- See the RESISTANCE MANAGEMENT section of the Coragen® eVo insect control Section 3 Federal label for additional guidance on insecticide resistance management.

**RESTRICTIONS AND PRECAUTIONS:**

- Do not apply more than 5.1 fl oz Coragen® eVo insect control or 0.2 lb ai of chlorantraniliprole containing products per acre per calendar year.
- Do not make more than 4 applications per acre per calendar year.
- Minimum interval between applications is 7 days.
- Preharvest interval is 14 days.
- The field re-entry interval is 4 hours.

R-4998 092925 09-29-25

FMC, the FMC logo and Coragen are Trademarks of FMC Corporation or an affiliate.  
©2025 FMC Corporation