CHLORANTRANILIPROLE	GROUP 28	INSECTICIDE

FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9661

FOR DISTRIBUTION ONLY IN THE STATE OF OREGON

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation for Coragen® eVo insect control is valid until December 31, 2031, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED SUBJECT TO THE TERMS AND CONDITIONS OF THE PRODUCT LABELING.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR CONTROL OF SQUASH VINE BORER IN CUCURBIT VEGETABLES

Directions For Use

CROP	APPLICATION METHOD	TARGET PEST	RATE	PHI (DAYS)	REI (HOURS)
Cucurbit Vegetables (EPA Crop Group 9)	Foliar	Squash Vine Borer (*Melittia cucurbitae*)	1.7 – 2.5 fl. oz./A (0.066 – 0.098 lb ai/A)	1	4

Cucurbit Vegetables (EPA Crop Group 9) Including: Chayote (fruit); Chinese wax- gourd (Chinese preserving melon); Citron melon; Cucumber; Gherkin; Edible gourd (includes hyotan, cucuzza, hechima, Chinese okra); Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); Muskmelon (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); Pumpkin, Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); Watermelon.

- Coragen® eVo insect control is recommended for the control of squash vine borer (*Melittia cucurbitae*) in cucurbit vegetables.
- Apply Coragen® eVo insect control at 1.7 fl. oz/A (0.066 lb ai/A) as a foliar spray for squash vine borer infestations in cucurbit vegetables. A non-ionic surfactant (NIS) is recommended.
- Spray programs should begin at flowering or at the first sign of squash vine borers.
- Make applications when populations reach local established treatment thresholds to prevent crop damage.
- Make repeat applications on a 7- to 10-day spray schedule if monitoring indicates the presence of squash vine borers.

FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000

Coragen, FMC and FMC logo – Trademarks of FMC or an affiliate
©2026 FMC Corporation

- See the RESISTANCE MANAGEMENT section of the Coragen® eVo insect control Section 3 Federal label for additional guidance on insecticide resistance management.
- Apply higher rates (up to 2.5 fl oz/A) for heavier infestations, larger/denser crops or extreme environmental conditions such as rainy weather and high temperatures.

USE RESTRICTIONS

- Do not make more than 4 applications per acre per crop or 12 applications per acre per calendar year.
- Minimum interval between treatments is 5 days for foliar applications and 10 days for drip chemigation applications.
- Do not apply more than 5.1 fl oz Coragen® eVo insect control or 0.2 lb a.i. of chlorantraniliprole-containing products per acre per crop.
- Do not apply more than 2.5 fl oz (0.098 lb ai per acre) Coragen® eVo insect control to the soil at planting.
- Do not apply more than 3.4 fl oz (0.132 lb ai per acre) Coragen® eVo insect control per crop by any combination of at plant soil application and drip chemigation.
- Do not make more than 2 drip chemigation applications of Coragen® eVo insect control per crop.
- Do not make more than one drip chemigation application per crop if an at plant application of Coragen® eVo insect control was made.
- Regardless of application method or chlorantraniliprole containing products used, do not apply more than 15.4 fl oz Coragen® eVo insect control or 0.6 lb ai of chlorantraniliprole-containing products per acre per calendar year. In NY do not apply more than 5.1 fl oz Coragen® eVo insect control or 0.2 lb ai of chlorantraniliprole- containing products per acre per calendar year.

IMPORTANT NOTICE

BEFORE BUYING OR USING FMC PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY, AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of Federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This Bulletin contains supplemental instructions for use of the referenced FMC product. Follow these instructions carefully. This Bulletin must be in the possession of the User at the time of application. **Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using the FMC product(s) covered by this Bulletin. FMC will not be responsible for losses or damages resulting from use of this product in any manner not specifically stated by FMC on the Section 3 label, or other labels, or this Bulletin. User assumes all risks associated with any such nonspecified use. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability of the Section 3 Federal product label.**

R-5055 072826 08-04-26

FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000

Coragen, FMC and FMC logo – Trademarks of FMC or an affiliate
©2026 FMC Corporation