(Container)

GROUP 4 HERBICIDE

FMC Fluroxypyr 333 EC Herbicide

FMC Fluroxypyr 333 EC is a selective herbicide for post-emergent control of annual broadleaved weeds including cleavers and certain ALS-resistant broadleaved weeds such as kochia in spring wheat, durum wheat, winter wheat, spring barley and oats. FMC Fluroxypyr 333 EC will also suppress hemp-nettle, common chickweedand wild buckwheat.

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: fluroxypyr, present as 1-methylheptyl ester......333 g/L

Emulsifiable concentrate

REGISTRATION NO. 35620 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1 L - 125 L

FMC of Canada Limited 6755 Mississauga Road, Suite 204 Mississauga, ON L5N 7Y2 1-833-362-7722

PRECAUTIONS KEEP OUT OF REACH OF CHILDREN

For agricultural/crop scenarios: **Do not** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12-hours.

Do not use in residential areas. Residential areas are defined as any use site where bystanders including children could be exposed during or after application. This includes homes, schools, parks, playgrounds, playing fields, public buildings, or any other area where the general public including children could be exposed.

Apply only when potential for drift to areas of human habitation and activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.

PROTECTIVE CLOTHING AND EQUIPMENT

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and work boots during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves are not required during application within a closed cab and/or cockpit.

For handheld application, wear eye, head and respiratory protection when applying above waist height, including overhead.

OPERATOR USE PRECAUTIONS

Wear freshly laundered clothing and clean protective equipment daily. At completion of spraying or end of the day take a shower immediately. Wash thoroughly with soap and water before eating, drinking, smoking or using the toilet. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse. If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Avoid contact with eyes, skin and clothing. Avoid breathing spray mist.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE. Do not use or store near heat or open flame.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

THIS PRODUCT CONTAINS A PETROLEUM DISTILLATE. Vomiting may cause aspiration pneumonia. The decision of whether to induce vomiting or not should be made by the attending physician. If lavage is performed, suggest endotracheal and/or oesophageal control. Danger from lung aspiration of petroleum based solvents must be weighed against toxicity when considering emptying the stomach. No specific antidote. Employ supportive care. Treatment should be based on the judgment of the physician in response to reactions of the patient.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

This product contains an active ingredient and aromatic petroleum distillates which are toxic to aquatic organisms.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

STORAGE

Store this product away from food or feed.

Store in original containers in a secure, dry heated storage. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL

Recyclable Containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

- 1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
- 2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable Containers

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

PMRA Approved Label – M.A. 2025-3103 2025-09-03

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, andfor clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

(Booklet)

GROUP 4 HERBICIDE

FMC Fluroxypyr 333 EC Herbicide

FMC Fluroxypyr 333 EC is a selective herbicide for post-emergent control of annual broadleaved weeds including cleavers and certain ALS-resistant broadleaved weeds such as kochia in spring wheat, durum wheat, winter wheat, spring barley and oats. FMC Fluroxypyr 333 EC will also suppress hemp-nettle, common chickweedand wild buckwheat.

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND BOOKLET BEFORE USING KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: fluroxypyr, present as 1-methylheptyl ester 333 g/L

Emulsifiable concentrate

REGISTRATION NO. 35620 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1 L - 125 L

FMC of Canada Limited 6755 Mississauga Road, Suite 204 Mississauga, ON L5N 7Y2 1-833-362-7722

PRECAUTIONS KEEP OUT OF REACH OF CHILDREN

For agricultural/crop scenarios: **Do not** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12-hours.

Do not use in residential areas. Residential areas are defined as any use site where bystanders including children could be exposed during or after application. This includes homes, schools, parks, playgrounds, playing fields, public buildings, or any other area where the general public including children could be exposed.

Apply only when potential for drift to areas of human habitation and activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.

PROTECTIVE CLOTHING AND EQUIPMENT

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and work boots during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves are not required during application within a closed cab and/or cockpit.

For handheld application, wear eye, head and respiratory protection when applying above waist height, including overhead.

OPERATOR USE PRECAUTIONS

Wear freshly laundered clothing and clean protective equipment daily. At completion of spraying or end of the day take a shower immediately. Wash thoroughly with soap and water before eating, drinking, smoking or using the toilet. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse. If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Avoid contact with eyes, skin and clothing. Avoid breathing spray mist.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE. Do not use or store near heat or open flame.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IN CASE OF EMERGENCY, CALL TOLL FREE, DAY OR NIGHT: 1-800-331-3148

THIS PRODUCT CONTAINS A PETROLEUM DISTILLATE. Vomiting may cause aspiration pneumonia. The decision of whether to induce vomiting or not should be made by the attending physician. If lavage is performed, suggest endotracheal and/or oesophageal control. Danger from lung aspiration of petroleum based solvents must be weighed against toxicity when considering emptying the stomach. No specific antidote. Employ supportive care. Treatment should be based on the judgment of the physician in response to reactions of the patient.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

This product contains an active ingredient and aromatic petroleum distillates which are toxic to aquatic organisms.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

STORAGE

Store away from food or feed.

Store in original containers in a secure, dry heated storage. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL

Recyclable Containers:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at acontainer collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

- 1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
- 2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable Containers:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the registrant or the provincial or

territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, andfor clean-up of spills.

GENERAL INFORMATION

FMC Fluroxypyr 333 EC is a selective post-emergent herbicide for the control of annual broadleaved weeds in spring wheat, durum wheat, winter wheat, spring barley and oats, not underseeded with legumes. FMC Fluroxypyr 333 EC offers a novel mode of action for controlling hard-to-kill annual broadleaved weeds such as kochia (including ALS resistant biotypes) and cleavers. FMC Fluroxypyr 333 EC will also provide suppression of hempnettle, wild buckwheat and chickweed.

MODE OF ACTION

FMC Fluroxypyr 333 EC is a systemic auxin-type herbicide and moves within the plant for control of exposed and underground plant tissues. The product controls weeds by disrupting normal plant growth patterns. Symptoms of weeds include epinasty (twisting of the stems) and swollen nodes.

GENERAL USE PRECAUTIONS

Sensitive Plants

Do not apply FMC Fluroxypyr 333 EC directly to, or otherwise permit it to come in direct contact with susceptible crops or desirable plants including alfalfa, edible beans, canola, flowers and ornamentals, lettuce, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tomatoes or tobacco.

Non-Target Sites

Do not apply where proximity of susceptible crops or other desirable plants is likely to result in exposure to spray or spray drift. See Environmental Hazards section of the label.

Crop Rotation

Fields previously treated with FMC Fluroxypyr 333 EC can be seeded the following year to alfalfa, barley, canola, corn, dry beans, fababeans, flax, forage grasses, lentils, mustard, oats, peas, potatoes, rye, soybeans, sugar beets, sunflowers or wheat or fields can be summerfallowed.

Tank Mixtures

This product may be tank mixed with a fertilizer, a supplement or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact FMC at 1-833-362-7722 for information before applying any tank mix that is not specifically recommended on this label.

Spray Equipment Precaution

Do not apply through any type of irrigation system.

To Reduce Spray Drift

- Use nozzles delivering higher volumes and coarser droplets.
- Use low pressures (200 to 275 kPa).
- Spot treatments should only be applied with a calibrated boom to prevent over-application.

Equipment Clean-Up

Equipment used to apply FMC Fluroxypyr 333 EC should not be used to apply other pesticides to sensitive crops without thorough cleaning. Contact your FMC Fluroxypyr 333 EC dealer for detailed equipment cleaning procedures.

DIRECTIONS FOR USE

READ THE ENTIRE LABEL BEFORE USE. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY

RESULT IN ERRATIC WEED CONTROL OR CROP DAMAGE. DO NOT APPLY TO CROPS UNDERSEEDED WITH LEGUMES.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Crops Registered

spring wheat durum wheat winter wheat spring barley oats

FMC FLUROXYPYR 333 EC ALONE

Apply FMC Fluroxypyr 333 EC with ground or aerial application equipment (see Application directions section) for post-emergent control of cleavers and kochia, including ALS resistant kochia and other broadleaved weeds infesting spring wheat, durum wheat, winter wheat, spring barley and oats. See following table for weeds controlled/suppressed and applications rates.

Weeds Controlled or Suppressed By FMC Fluroxypyr 333 EC Alone

Weeds Controlled	FMC Fluroxypyr 333 EC Rate (L/ha)
chickweed* (up to 8 cm)	0.41
cleavers (1-6 whorls)	0.21
cleavers (1-8 whorls)	0.31
kochia* (2-8 leaf)	0.31
round-leaved mallow (1-6 leaf)	0.41
volunteer flax (1-12 cm)	0.31
Weeds Suppressed	FMC Fluroxypyr 333 EC Rate (L/ha)
hempnettle (2-6 leaf)	0.41
stork's-bill (1-8 leaf)	0.31
wild buckwheat (1-8 leaf)	0.21

^{*} Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.

NOTE: FMC Fluroxypyr 333 EC activity is influenced by weather conditions. Optimum activity requires active crop and weed growth. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions, e.g., drought, heat or cold stress, or if weeds have initiated flowering, or if heavy infestations exist.

Application Timing

Apply to actively growing spring wheat, durum wheat, spring barley and oats from the 2 leaf crop growth stage up to and including initiation of stem elongation. Apply to winter wheat in the spring from the 3 tiller stage to just before the flag leaf stage. Apply when weeds are actively growing. Best results are obtained from applications made to seedling weeds. Only weeds emerged at the time of treatment will be controlled. Extreme growing conditions such as drought or near freezing temperature prior to, at and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Foliage that is wet at the time of application may decrease control. FMC Fluroxypyr 333 EC applications are rainfast within 1 hour after application. Under conditions of low crop and high weed density, control may be reduced.

Pre-harvest/Grazing Intervals (FMC Fluroxypyr 333 EC applied alone)

- Livestock may be grazed on treated crops 3 days following application.
- Do not harvest the treated crop within 60 days after application.

APPLICATION DIRECTIONS (GROUND AND AERIAL APPLICATIONS)

(1) Ground Application

Using ground equipment, apply FMC Fluroxypyr 333 EC alone or with tank mix partners as a broadcast treatment. Apply FMC Fluroxypyr 333 EC alone at the recommended rates in a spray volume of 50-100 L water/ha. When tank mixing, refer to the label of the tank mix partner label for additional instructions, water volumes, precautions and weeds controlled.

Field sprayer application

DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) medium classification. Boom height must be 60 cm or less above the crop or ground.

(2) Conventionally piloted aircraft Application

Do not use human flaggers.

Use FMC Fluroxypyr 333 EC alone or tank mixed with a product registered for aerial application as a broadcast treatment. Apply the recommended rates of FMC Fluroxypyr 333 EC and tank mix partner as indicated elsewhere on this label. Apply in a spray volume of 30-50 L water/ha. Refer to the tank mixpartner label for additional instructions, precautions and weeds controlled. DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan

Apply only by aircraft equipment, which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate specified for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate swath marking devices.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the registrant at 1-833-362-7722 or obtain technical advice from the distributor or your provincial or territorial agricultural representative. Application of this specific product must meet and/or conform to the following:

The interaction of many equipment-and-weather-related factors determine the potential for spray drift.
 Users are responsible for considering all these factors when making decisions.

The following drift management requirement must be followed to avoid off-target drift movement from aerial applications: Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

SPRAY BUFFER ZONES

A spray buffer zone is NOT required for

• uses with hand-held application equipment permitted on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts,woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

		Spray Buffer Zones (metres) Required for the Protection of:					
Method of application	Crop	Freshwater Habitat of Depths:		Estuarine/marine Habitat of Depths:		Terrestrial habitat	
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m		
Field sprayer	Spring wheat, durum wheat, winter wheat, spring barley and oats	1	0	1	1	3	

Aerial Spring Fixed Wing	5	0	1	1	95
--------------------------	---	---	---	---	----

whe duru	im Rotary wing	3	0	1	1	80
whe						
wint						
whe sprir						
barle						
	oats					

When tank mixes are permitted, consult the labels of the tank mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product for conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Mitigation portion of the Canada.ca website.

TANK MIX COMBINATION - FMC FLUROXYPYR 333 EC + 2,4-D ESTER HERBICIDE (Do not apply to oats)

For control of a wide spectrum of broadleaved weeds (see following table) apply FMC Fluroxypyr 333 EC tankmixed with 2,4-D ester herbicide. Apply post-emergence when weeds are in the seedling stage (2-4 leaf) and when spring wheat, durum wheat, winter wheat and spring barley are in the 4 leaf to flag leaf stage. Apply to winter wheat in the spring from the 3 tiller stage to just before the flag leaf stage. Application before the 4 leaf stage of wheat and barley may cause severe twisting of leaves and stem, and head deformities which may reduce yield up to 10%. Do not apply later than the flag leaf stage.

Rates of Applications

I. Low Rate: To control the following weeds apply a uniform spray containing 0.23 L/ha FMC Fluroxypyr 333 EC and 420 g a.e./ha 2,4-D ester herbicide (e.g., equivalent to 0.64 L/ha of 2,4-D Ester 700).

Weeds Controlled

bluebur goat's-beard shepherd's-purse burdock stinkweed hoary cress* canola (volunteer) sunflower (annual) kochia*** cleavers (1-6 whorls) vetch lamb's-quarters clovers (sweet) wild radish mustards (except green, dog & tansy) cocklebur wild mustard plantain flixweed wild buckwheat (1-6 leaf) prickly lettuce field horsetail* ragweed

II. High Rate: To control the above weeds plus the additional weeds listed below, apply a uniform spray containing 0.31 L/ha of FMC Fluroxypyr 333 EC and 560 g a.e./ha of 2,4-D ester herbicide (e.g., equivalent to 0.85 L/ha of 2,4-D Ester 700).

Additional Weeds Controlled:

blue lettuce* hairy galinsoga Russian thistle hedge bindweed smartweed cleavers (1-8 whorls) lady's-thumb stork's-bill (1-8 leaf) dandelion** tansy mustard leafy spurge* docks tartary buckwheat oak-leaved goosefoot dog mustard redroot pigweed volunteer flax (1-12 cm) field bindweed* wild buckwheat (1-8 leaf) round-leaved mallow (1-6 Leaf) field peppergrass

gumweed

Weeds Suppressed:

common chickweed*** (up to 8 cm) hempnettle (2-6 leaf stage) sow-thistle (perennial) * sow-thistle (annual)

- Top growth control only
- ** Spring rosettes
- *** Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme

NOTE: The activity of FMC Fluroxypyr 333 EC combined with 2,4-D ester herbicide is influenced by weather conditions. Optimum activity requires active crop and weed growth. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions, e.g., drought, heat or cold stress, or if weeds haveinitiated flowering, or if heavy infestations exist.

Mixing Instructions

Add the correct amount of FMC Fluroxypyr 333 EC to spray tank 1/2 filled with water and agitate. Add the correct amount of 2,4-D ester herbicide (e.g., 2,4-D Ester 700) and continue to agitate.

Pre-Harvest/Grazing Intervals (FMC Fluroxypyr 333 EC + 2,4-D ester herbicide)

- Do not permit lactating dairy animals to graze fields within 7 days after application.
- Do not harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.
- Do not harvest the treated mature crop within 60 days after application.

TANK MIX COMBINATIONS - FMC FLUROXYPYR 333 EC + 2,4-D ESTER HERBICIDE + OTHER HERBICIDES (Do not apply to oats)

Tank mixtures of FMC Fluroxypyr 333 EC + 2,4-D ester herbicide with other herbicides will provide control of additional weeds. For products which may be tank mixed with FMC Fluroxypyr 333 EC + 2,4-D ester herbicideand the additional weeds controlled see following table. Follow all precautions, minimal interval to harvestand directions for use on the FMC Fluroxypyr 333 EC and tank mix partner labels.

Tank Mix Combinations with FMC Fluroxypyr 333 EC (0.31 L/ha) + 2,4-D ester herbicide (560 g a.e./ha) or FMC Fluroxypyr 333 EC (0.23 L/ha) + 2,4-D ester herbicide (420 g a.e./ha)

Herbicide Tank Mix Partner	Crops Registered	Rate of Tank Mix Partner	Adjuvant Rate	Additional Weeds
Simplicity™◆◆◆	spring wheat, durum wheat, winter wheat	0.5 L/ha	None required	barnyard grass (1-5 leaf), common chickweed (up to 10 cm), hemp nettle (1-8 leaf), Japanese brome (1-6 leaf), wild oats (up to 4 leaf, 2 tiller), wild buckwheat (1-4 leaf)

^{**} Wild oat control may be reduced when tank mixed with FMC Fluroxypyr 333 EC + 2,4-D ester herbicide

TANK MIX COMBINATIONS - FMC FLUROXYPYR 333 EC + REFINE® SG HERBICIDE

For control of a wide spectrum of broadleaved weeds (see following table) FMC Fluroxypyr 333 EC can be tank mixed with Refine SG Herbicide. Apply post-emergence when weeds are actively growing, with best results on seedling weeds that are less than 10 cm in height. Only weeds emerged at the time of

^{***}Tank mixing with Simplicity only applies to the low rate of FMC Fluroxypyr 333 EC + 2,4-D ester

treatment will be controlled. Thorough covering of the target weeds is essential. The tankmix can be applied to actively growing spring wheat, including durum, spring barley and oats from the 2 leaf crop growth stage up to and including initiation of stem elongation. The tank mix of Refine SG Herbicide with FMC Fluroxypyr 333 EC allows for the control of Group 2 resistant cleavers at 0.21 L/ha and kochia 0.31 L/ha. Do not apply to crops underseeded to legumes or grasses.

Rates of Application To control the following weeds apply a uniform spray containing 0.21 to 0.41 litres of FMC Fluroxypyr 333 EC with 30 g/ha of Refine SG Herbicide and 2 litres per 1000 L spray solution of non-ionic surfactant (Agral 90*, Ag-Surf** or Citowett Plus) with a spray volume of 100 L/ha.

- * Choose one of Agral 90 Non-Ionic Wetting & Spreading Agent Liquid or Agral 90 Non-Ionic Liquid Wetting & Spreading Agent.
- ** Choose one of IPCO Ag-Surf Original, IPCO Ag-Surf II, Weedaway Ag-Surf Liquid Spray Adjuvant or Weedaway Ag-Surf II.

Weeds Controlled or Suppressed

Weeds Controlled at 0.21 L/ha of FMC Fluroxypyr 333 EC and 30 g/ha of Refine® SG Herbicide:

annual smartweed (green groundsel (common) shepherd's purse smartweed, lady's-thumb) hempnettle stinkweed ball mustard kochia² tartary buckwheat chickweed (1-6 leaf)1 lamb's-quarters volunteer rapeseed3 cleavers (1-4 whorls) narrow-leaved hawk's beard volunteer sunflower redroot pigweed wild buckwheat4 corn spurry cow cockle Russian thistle wild mustard flixweed

Weeds suppressed at 0.21 L/ha FMC Fluroxypyr 333 EC and 30 g/ha of Refine[®] SG Herbicide:

Canada thistle⁵ sow thistle⁵

round leaved mallow (2-6 leaf)⁶ stork's bill (2-6 leaf stage) scentless chamomile toadflax (up to 15 cm)⁷

Additional Weeds Controlled at 0.31 L/ha of FMC Fluroxypyr 333 EC and 30 g/ha of Refine[®] SG Herbicide:

kochia (2-8 leaf; including group 2 resistant biotypes) volunteer flax (1-12 cm)

Additional Weeds Controlled and Suppressed at 0.41 L/ha FMC Fluroxypyr 333 EC and 30 g/ha of Refine® SG Herbicide:

Weeds Controlled Weeds Suppressed

round leaved mallow (1-6 leaf) chickweed (up to 8 cm; including group 2 resistant biotypes)

¹ Spray before canopy prevents thorough coverage of weeds. Chickweed emerging after application willnot be controlled.

² Excluding Group 2 resistant kochia

³ The tank mix will not control imazethapyr tolerant canola varieties (e.g., Clearfield varieties or othervarieties with the Pursuit Smart trait)

⁴ Apply to actively growing wild buckwheat at the cotyledon to 3 leaf stage. Control may be reduced under environmental stress. Large plants may regrow after treatment.

⁵ Apply when most of the thistles have emerged and are actively growing. Apply before bud stage and before crop canopy prevents thorough coverage of thistles, when thistles are no larger than 15 cm in height for best top growth control. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be controlled.

⁶ Apply to actively growing round leaved mallow. Will keep mallow stunted, but may not be able to reduce overall populations

⁷ A control program for this weed includes chemical application and frequent tillage.

Mixing Instructions

Fill the tank to 1/3 full of clean water, and with the agitator running, add the required amount of Refine SG Herbicide and continue to agitate for a minimum of 5 minutes to ensure that Refine SG Herbicide is completely dissolved. Add the required amount of FMC Fluroxypyr 333 EC at a reduced agitation. Add the rest of the water and then add adjuvant. Use a spray volume of 100 L/ha.

Pre-Harvest/Grazing Intervals

- Livestock may not be grazed or fed crops within 7 days of application
- Do not harvest the treated crop within 60 days of application

TANK MIX COMBINATION - FMC FLUROXYPYR 333 EC + MCPA ESTER HERBICIDE

For control of a wide spectrum of broadleaved weeds (see following table) apply FMC Fluroxypyr 333 EC at 0.31 L/ha tank mixed with MCPA LV 500 Herbicide at 1.12 L/ha or MCPA Ester 600 Herbicide at 0.93 L/ha. Apply post-emergence when weeds are in the seedling stage (2-4 leaf) and when spring wheat, durum wheat, spring barley and oats are in the 3 leaf to flag leaf stage. Apply to winter wheat in the spring from the 3 tiller stage to just before the flag leaf stage. Application before the 3 leaf stage of wheat and barley may cause severe twisting of leaves and stem, and head deformities which may reduce yield up to 10%. Do not apply later than the flag leaf stage. Do not apply to cereals underseeded with a legume.

Mixing Instructions

Add the correct amount of FMC Fluroxypyr 333 EC to spray tank 1/2 filled with water and agitate. Add the correct amount of MCPA ester herbicide and continue to agitate.

Weeds Controlled or Suppressed by FMC Fluroxypyr 333 EC + MCPA ester herbicide Tank Mix Weeds Controlled

shepherd's-purse burdock, common kochia• lamb's-quarters canola, volunteer stinkweed cleavers (1-8 whorls) mustard (except dog & green tansy) sunflower, annual pigweed, redroot (1-4 leaf) cocklebur vetch flax, volunteer (1-12 cm) prickly lettuce wild radish flixweed ragweed, common wild buckwheat (1-8 leaf)

hempnettle (2-6 leaf)

Weeds Suppressed

green smartweed stork's-bill (1-8 leaf)

Pre-Harvest/Grazing Intervals (FMC Fluroxypyr 333 EC + MCPA ester herbicide)

- Allow 7 days between application and grazing/cutting for hay.
- Do not harvest the treated mature crop within 60 days after application.

TANK MIX COMBINATION - FMC FLUROXYPYR 333 EC +SIMPLICITY (Do not apply to oats)

For control of a wide spectrum of broadleaf and grass weeds (see following table) in spring wheat, durum wheat and winter wheat, apply FMC Fluroxypyr 333 EC tank mixed with Simplicity. Apply post-emergence when weeds are actively growing: grass weeds up to the 4 leaf, 2 tiller stage and broadleaf weeds as indicated in the following table. Best results are obtained from applications made to seedling weeds.

Apply to actively growing spring or durum wheat from the 3 leaf stage to the first node stage. Apply to winter wheat in the 2-7 leaf, 4 tiller stage with a spring application. Extreme growing conditions such as drought or near freezing temperature prior to, at, or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Under conditions of low crop and high weed

[•]Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.

density, control may be reduced.

Mixing Instructions

Fill the sprayer tank ½ full of water and start sprayer tank agitation. Add the required amount of Simplicity. Add the correct amount of FMC Fluroxypyr 333 EC and continue to agitate.

Pre-harvest/Grazing Intervals (FMC Fluroxypyr 333 EC and Simplicity)

- Livestock may be grazed on treated crops 7 days following application.
- Do not harvest the treated crop within 60 days after application.

Rates of Application

I. Low Rate: To control or suppress the following weeds apply a uniform spray containing FMC Fluroxypyr 333 EC at 0.21 L/ha tank mixed with Simplicity at 375 mL/ha.

Weeds Controlled

cleavers (1-6 whorls)

wild oats (up to 4 leaf, 2 tiller) *

*Under low wild oat populations (<75 plants/m²)

II. High Rate: To control the above weeds plus the additional weeds listed below, apply a uniform spray containing FMC Fluroxypyr 333 EC at 0.31 L/ha with Simplicity at 500 mL/ha.

Additional Weeds Controlled

barnyard grass (1-5 leaf)	kochia* (2-8 leaf)
chickweed, common (up to 10 cm)	red root pigweed (1-8 leaf)
cleavers (1-8 whorls)	smartweed (lady's thumb, 1-5 leaf)
foxtail, yellow (1-5 leaf)	volunteer canola * * (1-6 leaf)
hemp nettle (1-8 leaf)	volunteer flax (1-12 cm)
Japanese brome (1-6 leaf)	

^{*}Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.

Additional Weeds Suppressed

brome, downy (2-6 leaf, 2 tillers)	storks-bill (1-8 leaf)
foxtail, green (1-5 leaf)	wild buckwheat (1-4 leaf)

TANK MIX COMBINATIONS –FMC FLUROXYPYR 333 EC + SIMPLICITY + OTHER HERBICIDES (Do not apply to oats)

When tank mixing FMC Fluroxypyr 333 EC + Simplicity at either the low or high rates, the addition of other herbicides will provide control of additional weeds. For products which may be tank mixed with FMC Fluroxypyr 333 EC + Simplicity and the additional weeds controlled see following table. Follow all precautions, minimal interval to harvest and directions for use on the FMC Fluroxypyr 333 EC and tank mix partner labels.

Tank Mix Combinations with FMC Fluroxypyr 333 EC (0.21 L/ha) + Simplicity (375 mL/ha) or FMC Fluroxypyr 333 EC (0.31 L/ha) + Simplicity (500 mL/ha)

i luloxypyl 333 Lo (o.o i E/ila) · Olliplici	ty (300 iiiL/iia)	
Herbicide		Rate of Tank Mix	
Tank Mix Partner	Crops Registered	Partner	Additional Weeds Controlled

^{**} will not control volunteer Imidazolinone-tolerant canola

2,4-D ester*	spring wheat, durum wheat, winter wheat	Up to 420 g ae/ha	Susceptible weeds such as: mustard (except dog and green tansy), flixweed, bluebur, burdock, cocklebur, stinkweed, goatsbeard, prickly lettuce, lamb's-quarters, plantain, ragweeds, Russian thistle, shepherd's purse, annual sunflower, sweet clover, volunteer canola, wild buckwheat, wild radish *Consult 2,4-D ester labels for weed stages and rates
MCPA ester*	spring wheat durum wheat, winter wheat	Up to 560 g ae/ha	Burdock (before the 4 leaf stage), cocklebur, flixweed, plantain, lamb's-quarters, mustards (except dog and tansy), prickly lettuce, ragweeds, Russian pigweed, shepherd's-purse, stinkweed, vetch, wild buckwheat, wild radish, wild (annual) sunflower. *Consult MCPA ester labels for weed stages and rates

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, FMC Fluroxypyr 333 EC is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to FMC Fluroxypyr 333 EC and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the samefield. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of FMC Fluroxypyr 333 EC or other Group 4 herbicides with differentherbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To
 delay resistance, the less resistance-prone partner should control the target weed(s) as
 effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical informationrelated to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

PMRA Approved Label – M.A. 2025-3103 2025-09-03

 For further information or to report suspected resistance, contact FMC representatives at 1-833-362-7722.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

® REFINE are trademarks of FMC Corporation or an affiliate.

All other products listed are registered trademarks of their respective companies